Trigonometric Functions



Tidal Coastline and Pots of Water

In Florida, they post times of tides coming in and going out very precisely, like 11:23 A.M. How can they be so precise? There is more to tides, the rise and fall of ocean waters, than the gravitational pull of the Moon and Sun.

These are the chief factors, of course. And because the movements of Earth, Sun and Moon in relationship to each other are known precisely, the rhythm of tides rising and falling along coastlines is easy to predict.

Yet the time and heights of high and low tide may vary along different stretches of the same coast, though they are reacting to similar driving forces and pressures.

Historic observation makes possible the exact timing of high tides and low tides along a particular section of coast for a month, a year, or far into the future.

The reason for the difference is oscillation. Think of pots and pans filled with varying levels of water on a table, says Charles O' Reilly, Chief of Tidal Analysis for the Geological Survey of Canada's hydrographic service in Dartmouth, Nova Scotia. Then kick the table.

"You'll notice the water in the pots and pans will slosh differently. That's their natural oscillation," he said. "If you kick the table rhythmically, you'll find each pot continues to slosh differently because it has its own rhythm.

"Now, if you join those pots and pans together, that's sort of like a coastal ocean. They're all feeling the same 'kick,' but they are all responding differently. In order to predict a tide, you have to measure them for some period of time."

SOURCE: Toronto Star, June 13, 2001, p. GT02. Reprinted with permission—Torstar Syndication Services.

-See Chapter Project 1.



A LOOK BACK In Chapter 2, we began our discussion of functions. We defined domain and range and independent and dependent variables; we found the value of a function and graphed functions. We continued our study of functions by listing properties that a function might have, like being even or odd, and we created a library of functions, naming key functions and listing their properties, including the graph.

A LOOK AHEAD In this chapter we define the trigonometric functions, six functions that have a wide application. We shall talk about their domain and range, see how to find values and graph them, and develop a list of their properties.

There are two widely accepted approaches to the development of the trigonometric functions: one uses right triangles; the other uses circles, especially the unit circle. In this book, we develop the trigonometric functions using the unit circle. In Chapter 7, we present right triangle trigonometry.


OUTLINE

- 5.1 Angles and Their Measure
- **5.2** Trigonometric Functions: Unit Circle Approach
- 5.3 Properties of the Trigonometric Functions
- 5.4 Graphs of the Sine and Cosine Functions
- **5.5** Graphs of the Tangent, Cotangent, Cosecant, and Secant Functions
- 5.6 Phase Shift; Sinusoidal Curve Fitting Chapter Review Chapter Test Chapter Projects Cumulative Review

5.1 Angles and Their Measure					
Before getting started, review the following:					
(Appendix, Section A.2, p. 963)					
?' problems on page 366.					
Convert between Degrees, Minutes, Seconds, and Decimal Forms for Angles					
Find the Arc Length of a Circle					
Convert from Degrees to Radians and from Radians to Degrees					
Find the Area of a Sector of a Circle					
Find the Linear Speed of an Object Traveling in Circular Motion					


A ray, or half-line, is that portion of a line that starts at a point V on the line and extends indefinitely in one direction. The starting point V of a ray is called its vertex. See Figure 1.

If two rays are drawn with a common vertex, they form an **angle**. We call one of the rays of an angle the **initial side** and the other the **terminal side**. The angle formed is identified by showing the direction and amount of rotation from the initial side to the terminal side. If the rotation is in the counterclockwise direction, the angle is **positive**; if the rotation is clockwise, the angle is **negative**. See Figure 2. Lowercase Greek letters, such as α (alpha), β (beta), γ (gamma), and θ (theta), will be used to denote angles. Notice in Figure 2(a) that the angle α is positive because the direction of the rotation from the initial side to the terminal side is counterclockwise. The angle β in Figure 2(b) is negative because the rotation is clockwise. The angle γ in Figure 2(c) is positive. Notice that the angle α in Figure 2(a) and the angle γ in Figure 2(c) have the same initial side and the same terminal side. However, α and γ are unequal, because the amount of rotation required to go from the initial side to the terminal side is greater for angle γ than for angle α .




An angle θ is said to be in **standard position** if its vertex is at the origin of a rectangular coordinate system and its initial side coincides with the positive *x*-axis. See Figure 3.





When an angle θ is in standard position, the terminal side will lie either in a quadrant, in which case we say that θ lies in that quadrant, or θ will lie on the x-axis or the y-axis, in which case we say that θ is a quadrantal angle. For example, the angle θ in Figure 4(a) lies in quadrant II, the angle θ in Figure 4(b) lies in quadrant IV, and the angle θ in Figure 4(c) is a quadrantal angle.




We measure angles by determining the amount of rotation needed for the initial side to become coincident with the terminal side. The two commonly used measures for angles are *degrees* and *radians*.


Degrees

The angle formed by rotating the initial side exactly once in the counterclockwise direction until it coincides with itself (1 revolution) is said to measure 360 degrees, abbreviated 360°. **One degree, 1**°, is $\frac{1}{360}$ revolution. A **right angle** is an angle that measures 90°, or $\frac{1}{4}$ revolution; a **straight angle** is an angle that measures 180°, or $\frac{1}{2}$ revolution. See Figure 5. As Figure 5(b) shows, it is customary to indicate a right angle by using the symbol \square .





It is also customary to refer to an angle that measures θ degrees as an angle of θ degrees.



Convert between Degrees, Minutes, Seconds, and Decimal Forms for Angles

Although subdivisions of a degree may be obtained by using decimals, we also may

use the notion of *minutes* and *seconds*. **One minute**, denoted by **1**', is defined as $\frac{1}{60}$ degree. **One second**, denoted by **1**", is defined as $\frac{1}{60}$ minute, or equivalently, $\frac{1}{3600}$ degree. An angle of, say, 30 degrees, 40 minutes, 10 seconds is written compactly as $30^{\circ}40'10"$. To summarize:

1 counterclockwise revolution =
$$360^{\circ}$$

 $1^{\circ} = 60'$ $1' = 60''$ (1)

It is sometimes necessary to convert from the degree, minute, second notation $(D^{\circ}M'S'')$ to a decimal form, and vice versa. Check your calculator; it should be capable of doing the conversion for you.

Before getting started, though, you must set the mode to degrees because there are two common ways to measure angles: degree mode and radian mode. (We will define radians shortly.) Usually, a menu is used to change from one mode to

another. Check your owner's manual to find out how your particular calculator works.

Now let's see how to convert from the degree, minute, second notation $(D^{\circ}M'S'')$ to a decimal form, and vice versa, by looking at some examples:

$$15^{\circ}30' = 15.5^{\circ} \text{ because } 30' = 30 \cdot \left(\frac{1}{60}\right)^{\circ} = 0.5^{\circ}$$

$$1' = \left(\frac{1}{60}\right)^{\circ}$$

$$32.25^{\circ} = 32^{\circ}15' \text{ because } 0.25^{\circ} = \left(\frac{1}{4}\right)^{\circ} = \frac{1}{4}(60') = 15'$$

$$1^{\circ} = 60'$$

Converting between Degrees, Minutes, Seconds, EXAMPLE 2 and Decimal Forms

- (a) Convert $50^{\circ}6'21''$ to a decimal in degrees.
- (b) Convert 21.256° to the D°M'S" form.

Algebraic Solution

(a) Because $1' = \left(\frac{1}{60}\right)^{\circ}$ and $1'' = \left(\frac{1}{60}\right)' = \left(\frac{1}{60} \cdot \frac{1}{60}\right)^{\circ}$, we convert as follows:

$$50^{\circ}6'21'' = 50^{\circ} + 6' + 21''$$
$$= 50^{\circ} + 6 \cdot \left(\frac{1}{60}\right)^{\circ} + 21 \cdot \left(\frac{1}{60} \cdot \frac{1}{60}\right)^{\circ}$$
$$\approx 50^{\circ} + 0.1^{\circ} + 0.005833^{\circ}$$
$$= 50.105833^{\circ}$$

(b) We proceed as follows:

$$21.256^{\circ} = 21^{\circ} + 0.256^{\circ}$$

$$= 21^{\circ} + (0.256)(60')$$

$$= 21^{\circ} + 15.36'$$

$$= 21^{\circ} + 15' + 0.36'$$

$$= 21^{\circ} + 15' + (0.36)(60'')$$


$$= 21^{\circ} + 15' + 21.6''$$

$$\approx 21^{\circ}15'22''$$

Graphing Solution


(a) Figure 10 shows the solution using a TI-84 Plus graphing calculator.





(b) Figure 11 shows the solution using a TI-84 Plus graphing calculator.

Figure 11



NOW WORK PROBLEMS 23 AND 29.

to




In many applications, such as describing the exact location of a star or the precise position of a boat at sea, angles measured in degrees, minutes, and even seconds are used. For calculation purposes, these are transformed to decimal form. In other applications, especially those in calculus, angles are measured using *radians*.

Radians

A **central angle** is an angle whose vertex is at the center of a circle. The rays of a central angle subtend (intersect) an arc on the circle. If the radius of the circle is r and the length of the arc subtended by the central angle is also r, then the measure of the angle is **1 radian**. See Figure 12(a).

Figure 12



For a circle of radius 1, the rays of a central angle with measure 1 radian would subtend an arc of length 1. For a circle of radius 3, the rays of a central angle with measure 1 radian would subtend an arc of length 3. See Figure 12(b).

2 Find the Arc Length of a Circle

Now consider a circle of radius r and two central angles, θ and θ_1 , measured in radians. Suppose that these central angles subtend arcs of lengths s and s_1 , respectively, as shown in Figure 13. From geometry, we know that the ratio of the measures of the angles equals the ratio of the corresponding lengths of the arcs subtended by these angles; that is,

$$\frac{\theta}{\theta_1} = \frac{s}{s_1} \tag{2}$$

Suppose that $\theta_1 = 1$ radian. Refer again to Figure 12(a). The amount of arc s_1 subtended by the central angle $\theta_1 = 1$ radian equals the radius *r* of the circle. Then $s_1 = r$, so equation (2) reduces to


$$\frac{\theta}{1} = \frac{s}{r}$$
 or $s = r\theta$ (3)

Theorem

Arc Length

For a circle of radius r, a central angle of θ radians subtends an arc whose length s is

$$s = r\theta$$
 (4)



SECTION 5.1 Angles and Their Measure 361

NOTE Formulas must be consistent with regard to the units used. In equation (4), we write

 $s = r\theta$

To see the units, however, we must go back to equation (3) and write

$$\frac{\theta \text{ radians}}{1 \text{ radian}} = \frac{s \text{ length units}}{r \text{ length units}}$$
$$s \text{ length units} = r \text{ length units} \frac{\theta \text{ radians}}{1 \text{ radian}}$$

Since the radians cancel, we are left with

s length units = (r length units) θ $s = r\theta$

where θ appears to be "dimensionless" but, in fact, is measured in radians. So, in using the formula $s = r\theta$, the dimension for θ is radians, and any convenient unit of length (such as inches or meters) may be used for s and r.

EXAMPLE 3	Finding the Length of an Arc of a Circle
	Find the length of the arc of a circle of radius 2 meters subtended by a central angle of 0.25 radian.
Solution	We use equation (4) with $r = 2$ meters and $\theta = 0.25$. The length s of the arc is
	$s = r\theta = 2(0.25) = 0.5$ meter
	NOW WORK PROBLEM 71.

3 Convert from Degrees to Radians and from Radians to Degrees

Consider a circle of radius r. A central angle of 1 revolution will subtend an arc equal to the circumference of the circle (Figure 14). Because the circumference of a circle equals $2\pi r$, we use $s = 2\pi r$ in equation (4) to find that, for an angle θ of 1 revolution,

$$s = r\theta$$

$$2\pi r = r\theta$$

$$\theta = 1 \text{ revolution; } s = 2\pi r$$

$$\theta = 2\pi \text{ radians}$$

Solve for θ .

From this we have,

1 revolution =
$$2\pi$$
 radians (5)

Since 1 revolution = 360° , we have


 $360^\circ = 2\pi$ radians


or

$$180^\circ = \pi$$
 radians (6)

Divide both sides of equation (6) by 180. Then

1 degree =
$$\frac{\pi}{180}$$
 radian





Divide both sides of (6) by π . Then

$$\frac{180}{\pi}$$
 degrees = 1 radian

We have the following two conversion formulas:

1 degree =
$$\frac{\pi}{180}$$
 radian 1 radian = $\frac{180}{\pi}$ degrees (7)

EXAMPLE 4 Converting from Degrees to Radians

Convert each angle in degrees to radians.

(a) 60° (b) 150° (c) -45° (d) 90° (e) 107° **Solution** (a) $60^{\circ} = 60 \cdot 1 \text{ degree} = <math>60 \cdot \frac{\pi}{180} \text{ radian} = \frac{\pi}{3} \text{ radians}$ (b) $150^{\circ} = 150 \cdot \frac{\pi}{180} \text{ radian} = \frac{5\pi}{6} \text{ radians}$ (c) $-45^{\circ} = -45 \cdot \frac{\pi}{180} \text{ radian} = -\frac{\pi}{4} \text{ radian}$ (d) $90^{\circ} = 90 \cdot \frac{\pi}{180} \text{ radian} = \frac{\pi}{2} \text{ radians}$ (e) $107^{\circ} = 107 \cdot \frac{\pi}{180} \text{ radian} \approx 1.868 \text{ radians}$

Example 4 illustrates that angles that are fractions of a revolution, (a)–(d), are expressed in radian measure as fractional multiples of π , rather than as decimals. For example, a right angle, as in Example 4(d), is left in the form $\frac{\pi}{2}$ radians, which is exact, rather than using the approximation $\frac{\pi}{2} \approx \frac{3.1416}{2} = 1.5708$ radians.

NOW WORK PROBLEMS 35 AND 61.

EXAMPLE 5 Converting Radians to Degrees

Convert each angle in radians to degrees.

(a) $\frac{\pi}{6}$ radian (b) $\frac{3\pi}{2}$ radians (c) $-\frac{3\pi}{4}$ radians (d) $\frac{7\pi}{3}$ radians (e) 3 radians Solution (a) $\frac{\pi}{6}$ radian $= \frac{\pi}{6} \cdot 1$ radian $= \frac{\pi}{6} \cdot \frac{180}{\pi}$ degrees $= 30^{\circ}$

(b)
$$\frac{3\pi}{2}$$
 radians = $\frac{3\pi}{2} \cdot \frac{180}{\pi}$ degrees = 270°


(c)
$$-\frac{3\pi}{4}$$
 radians $= -\frac{3\pi}{4} \cdot \frac{180}{\pi}$ degrees $= -135^{\circ}$
(d) $\frac{7\pi}{3}$ radians $= \frac{7\pi}{3} \cdot \frac{180}{\pi}$ degrees $= 420^{\circ}$
(e) 3 radians $= 3 \cdot \frac{180}{\pi}$ degrees $\approx 171.89^{\circ}$

Table 1 lists the degree and radian measures of some commonly encountered angles. You should learn to feel equally comfortable using degree or radian measure for these angles.

Table 1	Degrees Radians	0° 0	30° <u>#</u> 6	45° <u>π</u> 4	$\frac{60^{\circ}}{\frac{\pi}{3}}$	90° <u>#</u> 2	$\frac{120^{\circ}}{\frac{2\pi}{3}}$	$\frac{135^{\circ}}{\frac{3\pi}{4}}$	150° <u>5π</u> 6	180° π
	Degrees		210° 7π	225° 5π	240° 4π	270° 3π	300° 5π	315° 7π	330° 11π	360°
	Radians		6	4	3	2	3	4	6	2π

EXAMPLE 6 Finding the Distance between Two Cities

See Figure 15(a). The latitude of a location L is the angle formed by a ray drawn from the center of Earth to the Equator and a ray drawn from the center of Earth to L. See Figure 15(b). Glasgow, Montana, is due north of Albuquerque, New Mexico. Find the distance between Glasgow (48°9′ north latitude) and Albuquerque (35°5′ north latitude). Assume that the radius of Earth is 3960 miles.



Solution The measure of the central angle between the two cities is $48^{\circ}9' - 35^{\circ}5' = 13^{\circ}4'$. We use equation (4), $s = r\theta$, but first we must convert the angle of $13^{\circ}4'$ to radians.

 $\theta = 13^{\circ}4' \approx 13.0667^{\circ} = 13.0667 \cdot \frac{\pi}{180}$ radian ≈ 0.228 radian

We use $\theta = 0.228$ radian and r = 3960 miles in equation (4). The distance between the two cities is

$$s = r\theta = 3960 \cdot 0.228 \approx 903$$
 miles

When an angle is measured in degrees, the degree symbol will always be shown. However, when an angle is measured in radians, we will follow the usual practice and omit the word *radians*. So, if the measure of an angle is given as $\frac{\pi}{6}$, it is understood to mean $\frac{\pi}{6}$ radian.

NOW WORK PROBLEM 101.

Find the Area of a Sector of a Circle

Consider a circle of radius r. Suppose that θ , measured in radians, is a central angle of this circle. See Figure 16. We seek a formula for the area A of the sector formed by the angle θ (shown in blue).

Now consider a circle of radius r and two central angles θ and θ_1 , both measured in radians. See Figure 17. From geometry, we know the ratio of the measures of the angles equals the ratio of the corresponding areas of the sectors formed by these angles. That is,

$$\frac{\theta}{\theta_1} = \frac{A}{A_1}$$

Suppose that $\theta_1 = 2\pi$ radians. Then A_1 = area of the circle = πr^2 . Solving for A, we find

$$A = A_1 \frac{\theta}{\theta_1} = \pi r^2 \frac{\theta}{2\pi} = \frac{1}{2} r^2 \theta$$

Area of a Sector

The area A of the sector of a circle of radius r formed by a central angle of θ radians is

$$A = \frac{1}{2}r^2\theta \tag{8}$$

EXAMPLE 7 Finding the Area of a Sector of a Circle


Find the area of the sector of a circle of radius 2 feet formed by an angle of 30° . Round the answer to two decimal places.


We use equation (8) with r = 2 feet and $\theta = 30^{\circ} = \frac{\pi}{6}$ radians. [Remember, in equa-**Solution** tion (8), θ must be in radians.] The area A of the sector is


$$A = \frac{1}{2}r^2\theta = \frac{1}{2}(2)^2\frac{\pi}{6} = \frac{\pi}{3}$$
 square feet ≈ 1.05 square feet

rounded to two decimal places.

NOW WORK PROBLEM 79.







5 Find the Linear Speed of an Object Traveling in Circular Motion

We have already defined the average speed of an object as the distance traveled divided by the elapsed time. Suppose that an object moves around a circle of radius r at a constant speed. If s is the distance traveled in time t around this circle, then the **linear speed** v of the object is defined as

$$v = \frac{s}{t}$$
(9)

As this object travels around the circle, suppose that θ (measured in radians) is the central angle swept out in time *t*. See Figure 18. Then the **angular speed** ω (the Greek letter omega) of this object is the angle (measured in radians) swept out, divided by the elapsed time; that is,

$$\omega = \frac{\theta}{t} \tag{10}$$

Angular speed is the way the turning rate of an engine is described. For example, an engine idling at 900 rpm (revolutions per minute) is one that rotates at an angular speed of

$$900 \frac{\text{revolutions}}{\text{minute}} = 900 \frac{\text{revolutions}}{\text{minute}} \cdot 2\pi \frac{\text{radians}}{\text{revolution}} = 1800\pi \frac{\text{radians}}{\text{minute}}$$

There is an important relationship between linear speed and angular speed:

linear speed =
$$v = \frac{s}{t} = \frac{r\theta}{t} = r\left(\frac{\theta}{t}\right)$$

(9) $g = r\theta$

Then, using equation (10), we obtain

 $v = r\omega \tag{11}$

where ω is measured in radians per unit time.


When using equation (11), remember that $v = \frac{s}{t}$ (the linear speed) has the dimensions of length per unit of time (such as feet per second or miles per hour), r (the radius of the circular motion) has the same length dimension as s, and ω (the angular speed) has the dimensions of radians per unit of time. If the angular speed is given in terms of *revolutions* per unit of time (as is often the case), be sure to convert it to *radians* per unit of time before attempting to use equation (11).


Finding Linear Speed

A child is spinning a rock at the end of a 2-foot rope at the rate of 180 revolutions per minute (rpm). Find the linear speed of the rock when it is released.

Solution Look at Figure 19. The rock is moving around a circle of radius r = 2 feet. The angular speed ω of the rock is

$$\omega = 180 \frac{\text{revolutions}}{\text{minute}} = 180 \frac{\text{revolutions}}{\text{minute}} \cdot 2\pi \frac{\text{radians}}{\text{revolution}} = 360\pi \frac{\text{radians}}{\text{minute}}$$





From equation (11), the linear speed v of the rock is

$$v = r\omega = 2 \text{ feet} \cdot 360\pi \frac{\text{radians}}{\text{minute}} = 720\pi \frac{\text{feet}}{\text{minute}} \approx 2262 \frac{\text{feet}}{\text{minute}}$$

The linear speed of the rock when it is released is 2262 ft/min \approx 25.7 mi/hr.

NOW WORK PROBLEM 97.

HISTORICAL FEATURE

Trigonometry was developed by Greek astronomers, who regarded the sky as the inside of a sphere, so it was natural that triangles on a sphere were investigated early (by Menelaus of Alexandria about AD 100) and that triangles in the plane were studied much later. The first book containing a systematic treatment of plane and spherical trigonometry was written by the Persian astronomer Nasir Eddin (about AD 1250).

Regiomontanus (1436–1476) is the person most responsible for moving trigonometry from astronomy into mathematics. His work was improved by Copernicus (1473–1543) and Copernicus's student Rhaeticus (1514–1576). Rhaeticus's book was the first to define the six trigonometric functions as ratios of sides of triangles, although he did not give the functions their present names. Credit for this is due to Thomas Finck (1583), but Finck's notation was by no means universally accepted at the time. The notation was finally stabilized by the textbooks of Leonhard Euler (1707–1783).

Trigonometry has since evolved from its use by surveyors, navigators, and engineers to present applications involving ocean tides, the rise and fall of food supplies in certain ecologies, brain wave patterns, and many other phenomena.

5.1 Assess Your Understanding

'Are You Prepared?'

Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

1. What is the formula for the circumference *C* of a circle of radius *r*? (p. 963)

Concepts and Vocabulary

- An angle θ is in _____ if its vertex is at the origin of a rectangular coordinate system and its initial side coincides with the positive x-axis.
- 4. On a circle of radius r, a central angle of θ radians subtends an arc of length s = ____; the area of the sector formed by this angle θ is A = ____.
- 5. An object travels around a circle of radius *r* with constant speed. If *s* is the distance traveled in time *t* around the circle and θ is the central angle (in radians) swept out in time *t*, then the linear speed of the object is $v = ___$ and the angular speed of the object is $\omega = ___$.

- 2. What is the formula for the area A of a circle of radius r? (p. 963)
- **6.** *True or False:* $\pi = 180$ *.*
- 7. True or False: $180^{\circ} = \pi$ radians.
- 8. *True or False:* On the unit circle, if s is the length of the arc subtended by a central angle θ , measured in radians, then $s = \theta$.
- 9. *True or False:* The area A of the sector of a circle of radius r formed by a central angle of θ degrees is $A = \frac{1}{2}r^2\theta$.
- **10.** *True or False:* For circular motion on a circle of radius *r*, linear speed equals angular speed divided by *r*.

Skill Building

In Problems 11–22, draw each angle.

11. 30°	12. 60°	13. 135°	14. −120°	15. 450°	16. 540°
17. $\frac{3\pi}{4}$	18. $\frac{4\pi}{3}$	19. $-\frac{\pi}{6}$	20. $-\frac{2\pi}{3}$	21. $\frac{16\pi}{3}$	22. $\frac{21\pi}{4}$

In Problems 23–28, convert each angle to a decimal in degrees. Round your answer to two decimal places. Verify your results using a graphing utility.					
23. 40°10′25″	24. 61°42′21″	25. 1°2′3″	26. 73°40′40″	27. 9°9′9″	28. 98°22′45″
In Problems 29–34,	convert each angle to L	℃M'S" form. Round y	our answer to the neares	st second. Verify your re	esults using a graphing utility.
29. 40.32°	30. 61.24°	31. 18.255°	32. 29.411°	33. 19.99°	34. 44.01°
In Problems 35–46	, convert each angle in	degrees to radians. Ex	press your answer as a	multiple of π .	
35. 30°	36. 120°	37. 240°	38. 330°	39. −60°	40. -30°
41. 180°	42. 270°	43. −135°	44. -225°	45. -90°	46. -180°
In Problems 47–58	, convert each angle in	radians to degrees.			
47. $\frac{\pi}{3}$	48. $\frac{5\pi}{6}$	49. $-\frac{5\pi}{4}$	50. $-\frac{2\pi}{3}$	51. $\frac{\pi}{2}$	52. 4π
53. $\frac{\pi}{12}$	54. $\frac{5\pi}{12}$	55. $-\frac{\pi}{2}$	56. – <i>π</i>	57. $-\frac{\pi}{6}$	58. $-\frac{3\pi}{4}$
In Problems 59–64	, convert each angle in	degrees to radians. Ex	press your answer in de	ecimal form, rounded	to two decimal places.
59. 17°	60. 73°	▶ 61. −40°	62. −51°	63. 125°	64. 350°
In Problems 65–70	, convert each angle in	radians to degrees. Ex	press your answer in de	ecimal form, rounded	to two decimal places.
65. 3.14	66. 0.75	67. 2	68. 3	69. 6.32	70. $\sqrt{2}$
In Problems 71–78, answers to three de		f the arc of a circle of r	adius r subtended by the	e central angle θ. Find	the missing quantity. Round
71. $r = 10$ meters		= ?	72. $r = 6$ feet,	$\theta = 2$ radians, $s =$	= ?
73. $\theta = \frac{1}{3}$ radian,	s = 2 feet, $r = ?$		74. $\theta = \frac{1}{4}$ radi	an, $s = 6$ centimete	rs, $r = ?$

3	4
75. $r = 5$ miles, $s = 3$ miles, $\theta = ?$	76. $r = 6$ meters, $s = 8$ meters, $\theta = ?$
77. $r = 2$ inches, $\theta = 30^{\circ}$, $s = ?$	78. $r = 3$ meters, $\theta = 120^{\circ}$, $s = ?$

In Problems 79–86, A denotes the area of the sector of a circle of radius r formed by the central angle θ . Find the missing quantity. Round answers to three decimal places.

79. $r = 10$ meters, $\theta = \frac{1}{2}$ radian, $A = ?$	80. $r = 6$ feet, $\theta = 2$ radians, $A = ?$
81. $\theta = \frac{1}{3}$ radian, $A = 2$ square feet, $r = ?$	82. $\theta = \frac{1}{4}$ radian, $A = 6$ square centimeters, $r = ?$
83. $r = 5$ miles, $A = 3$ square miles, $\theta = ?$	84. $r = 6$ meters, $A = 8$ square meters, $\theta = ?$
85. $r = 2$ inches, $\theta = 30^{\circ}$, $A = ?$	86. $r = 3$ meters, $\theta = 120^{\circ}$, $A = ?$

In Problems 87–90, find the length s and area A. Round answers to three decimal places.




Applications and Extensions

91. Minute Hand of a Clock The minute hand of a clock is 6 inches long. How far does the tip of the minute hand move in 15 minutes? How far does it move in 25 minutes?



- **92.** Movement of a Pendulum A pendulum swings through an angle of 20° each second. If the pendulum is 40 inches long, how far does its tip move each second?
- **93.** Area of a Sector Find the area of the sector of a circle of radius 4 meters formed by an angle of 45°. Round the answer to two decimal places.
- **94.** Area of a Sector Find the area of the sector of a circle of radius 3 centimeters formed by an angle of 60°. Round the answer to two decimal places.
- **95. Watering a Lawn** A water sprinkler sprays water over a distance of 30 feet while rotating through an angle of 135°. What area of lawn receives water?



- **96. Designing a Water Sprinkler** An engineer is asked to design a water sprinkler that will cover a field of 100 square yards that is in the shape of a sector of a circle of radius 50 yards. Through what angle should the sprinkler rotate?
- **97. Motion on a Circle** An object is traveling around a circle with a radius of 5 centimeters. If in 20 seconds a central


angle of $\frac{1}{3}$ radian is swept out, what is the angular speed of the object? What is its linear speed?

- **98.** Motion on a Circle An object is traveling around a circle with a radius of 2 meters. If in 20 seconds the object travels 5 meters, what is its angular speed? What is its linear speed?
- **99. Bicycle Wheels** The diameter of each wheel of a bicycle is 26 inches. If you are traveling at a speed of 35 miles per hour on this bicycle, through how many revolutions per minute are the wheels turning?



100. Car Wheels The radius of each wheel of a car is 15 inches. If the wheels are turning at the rate of 3 revolutions per second, how fast is the car moving? Express your answer in inches per second and in miles per hour.


In Problems 101–104, the latitude of a location L is the angle formed by a ray drawn from the center of Earth to the Equator and a ray drawn from the center of Earth to L. See the figure.




- **101. Distance between Cities** Memphis, Tennessee, is due north of New Orleans, Louisiana. Find the distance between Memphis (35°9' north latitude) and New Orleans (29°57' north latitude). Assume that the radius of Earth is 3960 miles.
 - **102.** Distance between Cities Charleston, West Virginia, is due north of Jacksonville, Florida. Find the distance between Charleston (38°21' north latitude) and Jacksonville (30°20' north latitude). Assume that the radius of Earth is 3960 miles.
 - **103. Linear Speed on Earth** Earth rotates on an axis through its poles. The distance from the axis to a location on Earth 30° north latitude is about 3429.5 miles. Therefore, a location on Earth at 30° north latitude is spinning on a circle of radius 3429.5 miles. Compute the linear speed on the surface of Earth at 30° north latitude.
 - **104.** Linear Speed on Earth Earth rotates on an axis through its poles. The distance from the axis to a location on Earth 40° north latitude is about 3033.5 miles. Therefore, a location on Earth at 40° north latitude is spinning on a circle of radius 3033.5 miles. Compute the linear speed on the surface of Earth at 40° north latitude.
 - 105. Speed of the Moon The mean distance of the Moon from Earth is 2.39×10^5 miles. Assuming that the orbit of the Moon around Earth is circular and that 1 revolution takes 27.3 days, find the linear speed of the Moon. Express your answer in miles per hour.
 - **106. Speed of Earth** The mean distance of Earth from the Sun is 9.29×10^7 miles. Assuming that the orbit of Earth around the Sun is circular and that 1 revolution takes 365 days, find the linear speed of Earth. Express your answer in miles per hour.
 - **107. Pulleys** Two pulleys, one with radius 2 inches and the other with radius 8 inches, are connected by a belt. (See the figure.) If the 2-inch pulley is caused to rotate at 3 revolu-

tions per minute, determine the revolutions per minute of the 8-inch pulley.

[**Hint:** The linear speeds of the pulleys are the same; both equal the speed of the belt.]




- **108.** Ferris Wheels A neighborhood carnival has a Ferris wheel whose radius is 30 feet. You measure the time it takes for one revolution to be 70 seconds. What is the linear speed (in feet per second) of this Ferris wheel? What is the angular speed in radians per second?
- **109. Computing the Speed of a River Current** To approximate the speed of the current of a river, a circular paddle wheel with radius 4 feet is lowered into the water. If the current causes the wheel to rotate at a speed of 10 revolutions per minute, what is the speed of the current? Express your answer in miles per hour.



- **110. Spin Balancing Tires** A spin balancer rotates the wheel of a car at 480 revolutions per minute. If the diameter of the wheel is 26 inches, what road speed is being tested? Express your answer in miles per hour. At how many revolutions per minute should the balancer be set to test a road speed of 80 miles per hour?
- **111. The Cable Cars of San Francisco** At the Cable Car Museum you can see the four cable lines that are used to pull cable cars up and down the hills of San Francisco. Each cable travels at a speed of 9.55 miles per hour, caused by a rotating wheel whose diameter is 8.5 feet. How fast is the wheel rotating? Express your answer in revolutions per minute.
- **112.** Difference in Time of Sunrise Naples, Florida, is approximately 90 miles due west of Ft. Lauderdale. How much sooner would a person in Ft. Lauderdale first see the rising Sun than a person in Naples?


[**Hint:** Consult the figure. When a person at Q sees the first rays of the Sun, a person at P is still in the dark. The person at P sees the first rays after Earth has rotated so that P is at the location Q. Now use the fact that at the latitude of Ft. Lauderdale in 24 hours a length of arc of $2\pi(3559)$ miles is subtended.]



- **113. Keeping Up with the Sun** How fast would you have to travel on the surface of Earth at the equator to keep up with the Sun (that is, so that the Sun would appear to remain in the same position in the sky)?
- **114.** Nautical Miles A nautical mile equals the length of arc subtended by a central angle of 1 minute on a great circle^{*} on the surface of Earth. (See the figure.) If the radius of Earth is taken as 3960 miles, express 1 nautical mile in terms of ordinary, or statute, miles.



115. Approximating the Circumference of Earth Eratosthenes of Cyrene (276–194 BC) was a Greek scholar who lived and worked in Cyrene and Alexandria. One day while visiting in Syene he noticed that the Sun's rays shone directly down a well. On this date 1 year later, in Alexandria, which is 500 miles due north of Syene he measured the angle of the Sun to be about 7.2 degrees. See the figure. Use this information to approximate the radius and circumference of Earth.



^{*}Any circle drawn on the surface of Earth that divides Earth into two equal hemispheres.

370 CHAPTER 5 Trigonometric Functions

116. Pulleys Two pulleys, one with radius r_1 and the other with radius r_2 , are connected by a belt. The pulley with radius r_1 rotates at ω_1 revolutions per minute, whereas the pulley

Discussion and Writing

- **117.** Do you prefer to measure angles using degrees or radians? Provide justification and a rationale for your choice.
- **118.** What is 1 radian?
- **119.** Which angle has the larger measure: 1 degree or 1 radian? Or are they equal?
- **120.** Explain the difference between linear speed and angular speed.
- **121.** For a circle of radius *r*, a central angle of θ degrees subtends an arc whose length *s* is $s = \frac{\pi}{180}r\theta$. Discuss whether this is a

'Are You Prepared?' Answers

1. $C = 2\pi r$ **2.** $A = \pi r^2$

with radius r_2 rotates at ω_2 revolutions per minute. Show that $\frac{r_1}{r_2} = \frac{\omega_2}{\omega_1}$.

true or false statement. Give reasons to defend your position.

- **122.** Discuss why ships and airplanes use nautical miles to measure distance. Explain the difference between a nautical mile and a statute mile.
- **123.** Investigate the way that speed bicycles work. In particular, explain the differences and similarities between 5-speed and 9-speed derailleurs. Be sure to include a discussion of linear speed and angular speed.
- **124.** In Example 6, we found that the distance between Albuquerque, NM and Glasgow, MT is approximately 903 miles. According to mapquest.com, the distance is approximately 1300 miles. What might account for the difference?

5.2 Trigonometric Functions: Unit Circle Approach

PREPARING FOR THIS SECTION Before getting started, review the following:

- Pythagorean Theorem (Appendix, Section A.2, pp. 961–962)
- Unit Circle (Section 1.5, p. 45)

- Symmetry (Section 1.2, pp. 17–19)
- Functions (Section 2.1, pp. 56–65)

Now work the 'Are You Prepared?' problems on page 384.

OBJECTIVES 1 Find the Exact Values of the Trigonometric Functions Using a Point on the Unit Circle

- 2 Find the Exact Values of the Trigonometric Functions of Quadrantal Angles
- ³ Find the Exact Values of the Trigonometric Functions of $\frac{\pi}{4} = 45^{\circ}$
- 4 Find the Exact Values of the Trigonometric Functions of $\frac{\pi}{4} = 30^{\circ}$ and $\frac{\pi}{3} = 60^{\circ}$
- 5 Find the Exact Values of the Trigonometric Functions for Integer Multiples of

$$\frac{\pi}{6} = 30^{\circ}, \frac{\pi}{4} = 45^{\circ}, \text{ and } \frac{\pi}{3} = 60^{\circ}$$

- 6 Use a Calculator to Approximate the Value of a Trigonometric Function
- 7 Use a Circle of Radius r to Evaluate the Trigonometric Functions

We are now ready to introduce trigonometric functions. The approach that we take uses the unit circle.

The Unit Circle

Recall that the unit circle is a circle whose radius is 1 and whose center is at the origin of a rectangular coordinate system. Also recall that any circle of radius r has circumference of length $2\pi r$. Therefore, the unit circle (radius = 1) has a circumference of length 2π . In other words, for 1 revolution around the unit circle the length of the arc is 2π units.

The following discussion sets the stage for defining the trigonometric functions using the unit circle.


Let t be any real number. We position the t-axis so it is vertical with the positive direction up. We place this t-axis in the xy-plane, so that t = 0 is located at the point (1, 0) in the xy-plane.

If $t \ge 0$, let *s* be the distance from the origin to *t* on the *t*-axis. See the red portion of Figure 20(a).

Now look at the unit circle in Figure 20(a). Beginning at the point (1, 0) on the unit circle, travel s = t units in the counterclockwise direction along the circle, to arrive at the point P = (x, y). In this sense, the length s = t units is being **wrapped** around the unit circle.

If t < 0, we begin at the point (1, 0) on the unit circle and travel s = |t| units in the clockwise direction to arrive at the point P = (x, y). See Figure 20(b).





If $t > 2\pi$ or if $t < -2\pi$, it will be necessary to travel around the unit circle more than once before arriving at the point *P*. Do you see why?

Let's describe this process another way. Picture a string of length s = |t| units being wrapped around a circle of radius 1 unit. We start wrapping the string around the circle at the point (1, 0). If $t \ge 0$, we wrap the string in the counterclockwise direction; if t < 0, we wrap the string in the clockwise direction. The point P = (x, y)is the point where the string ends.

This discussion tells us that, for any real number t, we can locate a unique point P = (x, y) on the unit circle. We call P **the point on the unit circle that corresponds to t.** This is the important idea here. No matter what real number t is chosen, there is a unique point P on the unit circle corresponding to it. We use the coordinates of the point P = (x, y) on the unit circle corresponding to the real number t to define the **six trigonometric functions of** t.

Let t be a real number and let P = (x, y) be the point on the unit circle that corresponds to t.

The sine function associates with t the y-coordinate of P and is denoted by

 $\sin t = y$

The **cosine function** associates with *t* the *x*-coordinate of *P* and is denoted by

 $\cos t = x$

If $x \neq 0$, the **tangent function** is defined as

$\tan t = \frac{y}{2}$	
x	

If $y \neq 0$, the **cosecant function** is defined as

 $\csc t = \frac{1}{y}$

If $x \neq 0$, the secant function is defined as

 $\sec t = \frac{1}{x}$

If $y \neq 0$, the **cotangent function** is defined as

 $\cot t = \frac{x}{y}$

Notice in these definitions that if x = 0, that is, if the point P is on the y-axis, then the tangent function and the secant function are undefined. Also, if y = 0, that is, if the point P is on the x-axis, then the cosecant function and the cotangent function are undefined.

Because we use the unit circle in these definitions of the trigonometric functions, they are also sometimes referred to as **circular functions**.


Find the Exact Values of the Trigonometric Functions Using a Point on the Unit Circle

EXAMPLE 1Finding the Values of the Six Trigonometric Functions Using a
Point on the Unit Circle

Let t be a real number and let $P = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ be the point on the unit circle that corresponds to t. Find the values of sin t, cos t, tan t, csc t, sec t, and cot t.

2

Figure 21



Solution See Figure 21. We follow the definition of the six trigonometric functions, using $P = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right) = (x, y)$. Then, with $x = -\frac{1}{2}, y = \frac{\sqrt{3}}{2}$, we have $\cos t = x = -\frac{1}{2}$ $\tan t = \frac{y}{x} = \frac{\sqrt{3}}{\frac{2}{1}} = -\sqrt{3}$ $\sin t = v = \frac{\sqrt{3}}{2}$

$$csc t = \frac{1}{y} = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2\sqrt{3}}{3} \qquad sec t = \frac{1}{x} = \frac{1}{-\frac{1}{2}} = -2 \qquad cot t = \frac{x}{y} = \frac{-\frac{1}{2}}{\frac{\sqrt{3}}{2}} = -\frac{\sqrt{3}}{3}$$

NOW WORK PROBLEM 11.

2


Trigonometric Functions of Angles

Let P = (x, y) be the point on the unit circle corresponding to the real number t. See Figure 22(a). Let θ be the angle in standard position, measured in radians, whose terminal side is the ray from the origin through P. See Figure 22(b). Since the unit circle has radius 1 unit, from the formula for arc length, $s = r\theta$, we find that



So, if s = |t| units, then $\theta = t$ radians. See Figures 22(c) and (d).

Figure 22



The point P = (x, y) on the unit circle that corresponds to the real number t is the point P on the terminal side of the angle $\theta = t$ radians. As a result, we can say that

> $\sin t = \sin \theta$ $\begin{array}{c}
> \uparrow \\
> \text{Real number} \\
> \theta = t \text{ radians}
> \end{array}$

and so on. We can now define the trigonometric functions of the angle θ .

If $\theta = t$ radians, the six trigonometric functions of the angle θ are defined as

 $\sin \theta = \sin t$ $\cos \theta = \cos t$ $\tan \theta = \tan t$ $\csc \theta = \csc t$ $\sec \theta = \sec t$ $\cot \theta = \cot t$ Even though the distinction between trigonometric functions of real numbers and trigonometric functions of angles is important, it is customary to refer to trigonometric functions of real numbers and trigonometric functions of angles collectively as the *trigonometric functions*. We shall follow this practice from now on.

If an angle θ is measured in degrees, we shall use the degree symbol when writing a trigonometric function of θ , as, for example, in sin 30° and tan 45°. If an angle θ is measured in radians, then no symbol is used when writing a trigonometric function of θ , as, for example, in cos π and sec $\frac{\pi}{2}$.

Finally, since the values of the trigonometric functions of an angle θ are determined by the coordinates of the point P = (x, y) on the unit circle corresponding to θ , the units used to measure the angle θ are irrelevant. For example, it does not matter whether we write $\theta = \frac{\pi}{2}$ radians or $\theta = 90^{\circ}$. The point on the unit circle corresponding to this angle is P = (0, 1). As a result,

$$\sin \frac{\pi}{2} = \sin 90^\circ = 1$$
 and $\cos \frac{\pi}{2} = \cos 90^\circ = 0$

Find the Exact Values of the Trigonometric Functions of Quadrantal Angles

To find the exact value of a trigonometric function of an angle θ or a real number t requires that we locate the point P = (x, y) on the unit circle that corresponds to t. This is not always easy to do. In the examples that follow, we will evaluate the trigonometric functions of certain angles or real numbers for which this process is relatively easy. A calculator will be used to evaluate the trigonometric functions of most other angles.

EXAMPLE 2 Finding the Exact Values of the Six Trigonometric Functions of Quadrantal Angles

Find the exact values of the six trigonometric functions of:

(a)
$$\theta = 0 = 0^{\circ}$$
 (b) $\theta = \frac{\pi}{2} = 90^{\circ}$
(c) $\theta = \pi = 180^{\circ}$ (d) $\theta = \frac{3\pi}{2} = 270^{\circ}$

Solution (a) The point on the unit circle that corresponds to $\theta = 0 = 0^{\circ}$ is P = (1, 0). See Figure 23(a). Then

$$\sin 0 = \sin 0^\circ = y = 0$$
 $\cos 0 = \cos 0^\circ = x = 1$
 $\tan 0 = \tan 0^\circ = \frac{y}{x} = 0$ $\sec 0 = \sec 0^\circ = \frac{1}{x} = 1$

Since the y-coordinate of P is 0, csc 0 and cot 0 are not defined.




Figure 23






Figure 23







(b) The point on the unit circle that corresponds to $\theta = \frac{\pi}{2} = 90^{\circ}$ is P = (0, 1). See Figure 23(b). Then

$$\sin \frac{\pi}{2} = \sin 90^\circ = y = 1 \qquad \cos \frac{\pi}{2} = \cos 90^\circ = x = 0$$
$$\csc \frac{\pi}{2} = \csc 90^\circ = \frac{1}{y} = 1 \qquad \cot \frac{\pi}{2} = \cot 90^\circ = \frac{x}{y} = 0$$

Since the *x*-coordinate of *P* is 0, $\tan \frac{\pi}{2}$ and $\sec \frac{\pi}{2}$ are not defined.

(c) The point on the unit circle that corresponds to $\theta = \pi = 180^{\circ}$ is P = (-1, 0). See Figure 23(c). Then

 $\sin \pi = \sin 180^\circ = y = 0$ $\cos \pi = \cos 180^\circ = x = -1$

$$\tan \pi = \tan 180^\circ = \frac{y}{x} = 0$$
 $\sec \pi = \sec 180^\circ = \frac{1}{x} = -1$

Since the y-coordinate of P is 0, $\csc \pi$ and $\cot \pi$ are not defined.

(d) The point on the unit circle that corresponds to $\theta = \frac{3\pi}{2} = 270^{\circ}$ is P = (0, -1). See Figure 23(d). Then

$$\sin \frac{3\pi}{2} = \sin 270^\circ = y = -1 \qquad \cos \frac{3\pi}{2} = \cos 270^\circ = x = 0$$
$$\csc \frac{3\pi}{2} = \csc 270^\circ = \frac{1}{y} = -1 \qquad \cot \frac{3\pi}{2} = \cot 270^\circ = \frac{x}{y} = 0$$

Since the *x*-coordinate of *P* is 0, $\tan \frac{3\pi}{2}$ and $\sec \frac{3\pi}{2}$ are not defined.

Table 2 summarizes the values of the trigonometric functions found in Example 2.

Table 2

			Quadran	tal Angles			
θ (Radians)	heta (Degrees)	$\sin heta$	$\cos \theta$	tan $ heta$	$\csc \theta$	$\sec \theta$	$\cot heta$
0	0°	0	1	0	Not defined	1	Not defined
$\frac{\pi}{2}$	90°	1	0	Not defined	1	Not defined	0
π	180°	0	-1	0	Not defined	-1	Not defined
$\frac{3\pi}{2}$	270°	-1	0	Not defined	-1	Not defined	0


There is no need to memorize Table 2. To find the value of a trigonometric function of a quadrantal angle, draw the angle and apply the definition, as we did in Example 2.

EXAMPLE 3 Finding Exact Values of the Trigonometric Functions of Angles That Are Integer Multiples of Quadrantal Angles Find the exact value of: (a) $\sin(3\pi)$ (b) $\cos(-270^{\circ})$ **Solution** (a) See Figure 24. The point P on (b) See Figure 25. The point P on the unit circle that the unit circle that corresponds to $\theta = -270^{\circ}$ is corresponds to $\theta = 3\pi$ is P = (-1, 0), so $\sin(3\pi) = 0$. P = (0, 1), so $\cos(-270^{\circ}) = 0$. Figure 24 Figure 25 P = (0, 1)P = (-1, 0)NOW WORK PROBLEMS 19 AND 63. Find the Exact Values of the Trigonometric Functions of $\frac{\pi}{4} = 45^{\circ}$

EXAMPLE 4 Finding the Exact Values of the Trigonometric Functions of $\frac{\pi}{4} = 45^{\circ}$

Find the exact values of the six trigonometric functions of $\frac{\pi}{4} = 45^{\circ}$.

Solution We seek the coordinates of the point P = (x, y) on the unit circle that corresponds to $\theta = \frac{\pi}{4} = 45^{\circ}$. See Figure 26. First, we observe that *P* lies on the line y = x. (Do you see why? Since $\theta = 45^{\circ} = \frac{1}{2} \cdot 90^{\circ}$, *P* must lie on the line that bisects quadrant I.) Since P = (x, y) also lies on the unit circle, $x^2 + y^2 = 1$, it follows that



$$x^{2} + y^{2} = 1$$

$$x^{2} + x^{2} = 1 y = x, x > 0, y > 0$$

$$2x^{2} = 1$$

$$x = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}, y = \frac{\sqrt{2}}{2}$$

Then

$$\sin \frac{\pi}{4} = \sin 45^{\circ} = \frac{\sqrt{2}}{2} \qquad \cos \frac{\pi}{4} = \cos 45^{\circ} = \frac{\sqrt{2}}{2} \qquad \tan \frac{\pi}{4} = \tan 45^{\circ} = \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = 1$$

$$\csc \frac{\pi}{4} = \csc 45^{\circ} = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2} \qquad \sec \frac{\pi}{4} = \sec 45^{\circ} = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2} \qquad \cot \frac{\pi}{4} = \cot 45^{\circ} = \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = 1$$

EXAMPLE 5 Finding the Exact Value of a Trigonometric Expression


Find the exact value of each expression.

(a)
$$\sin 45^{\circ} \cos 180^{\circ}$$
 (b) $\tan \frac{\pi}{4} - \sin \frac{3\pi}{2}$ (c) $\left(\sec \frac{\pi}{4}\right)^{2} + \csc \frac{\pi}{2}$
Solution (a) $\sin 45^{\circ} \cos 180^{\circ} = \frac{\sqrt{2}}{2} \cdot (-1) = -\frac{\sqrt{2}}{2}$
From Example 4 From Table 2
(b) $\tan \frac{\pi}{4} - \sin \frac{3\pi}{2} = 1 - (-1) = 2$
From Example 4 From Table 2
(c) $\left(\sec \frac{\pi}{4}\right)^{2} + \csc \frac{\pi}{2} = (\sqrt{2})^{2} + 1 = 2 + 1 = 3$


Find the Exact Values of the Trigonometric Functions of $\frac{\pi}{6} = 30^{\circ}$ and $\frac{\pi}{3} = 60^{\circ}$

Consider a right triangle in which one of the angles is $\frac{\pi}{6} = 30^{\circ}$. It then follows that the third angle is $\frac{\pi}{3} = 60^{\circ}$. Figure 27(a) illustrates such a triangle with hypotenuse of length 1. Our problem is to determine *a* and *b*.

We begin by placing next to this triangle another triangle congruent to the first, as shown in Figure 27(b). Notice that we now have a triangle whose angles are each 60° . This triangle is therefore equilateral, so each side is of length 1. In particular,



the base is 2a = 1, and so $a = \frac{1}{2}$. By the Pythagorean Theorem, b satisfies the equation $a^2 + b^2 = c^2$, so we have




This results in Figure 27(c).

Finding the Exact Values of the Trigonometric Functions **EXAMPLE 6** of $\frac{\pi}{3} = 60^\circ$

Find the exact values of the six trigonometric functions of $\frac{\pi}{3} = 60^{\circ}$.

Solution Position the triangle in Figure 27(c) so that the 60° angle is in the standard position. See Figure 28.

Figure 28



The point on the unit circle that corresponds to $\theta = \frac{\pi}{3} = 60^{\circ}$ is $P = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. Then

$$\sin\frac{\pi}{3} = \sin 60^\circ = \frac{\sqrt{3}}{2} \qquad \qquad \cos\frac{\pi}{3} = \cos 60^\circ = \frac{1}{2}$$
$$\csc\frac{\pi}{3} = \csc 60^\circ = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2\sqrt{3}}{\sqrt{3}} \qquad \sec\frac{\pi}{3} = \sec 60^\circ = \frac{1}{\frac{1}{2}} = 2$$
$$\tan\frac{\pi}{3} = \tan 60^\circ = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \sqrt{3} \qquad \qquad \cot\frac{\pi}{3} = \cot 60^\circ = \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$$

EXAMPLE 7 Finding the Exact Values of the Trigonometric Functions of $\frac{\pi}{6} = 30^{\circ}$

Find the exact values of the trigonometric functions of $\frac{\pi}{6} = 30^{\circ}$.

Solution

 $P=(x, y)=\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$

<u>√3</u>

Position the triangle in Figure 27(c) so that the 30° angle is in the standard position. See Figure 29. π

The point on the unit circle that corresponds to $\theta = \frac{\pi}{6} = 30^{\circ}$ is $P = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$. Then

$$\sin\frac{\pi}{6} = \sin 30^{\circ} = \frac{1}{2} \qquad \qquad \cos\frac{\pi}{6} = \cos 30^{\circ} = \frac{\sqrt{3}}{2} \\ \csc\frac{\pi}{6} = \csc 30^{\circ} = \frac{1}{\frac{1}{2}} = 2 \qquad \qquad \sec\frac{\pi}{6} = \sec 30^{\circ} = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

$$\tan\frac{\pi}{6} = \tan 30^\circ = \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3} \qquad \cot\frac{\pi}{6} = \cot 30^\circ = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \sqrt{3}$$

Table 3 summarizes the information just derived for $\frac{\pi}{6} = 30^{\circ}, \frac{\pi}{4} = 45^{\circ}$, and $\frac{\pi}{3} = 60^{\circ}$. Until you memorize the entries in Table 3, you should draw an appropriate diagram to determine the values given in the table.

Table 3	θ (Radians)	θ (Degrees)	$\sin heta$	$\cos \theta$	tan θ	$\csc \theta$	$\sec \theta$	$\cot \theta$
	$\frac{\pi}{6}$	30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
	$\frac{\pi}{4}$	45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
	$\frac{\pi}{3}$	60°	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$

NOW WORK PROBLEM 39.

Constructing a Rain Gutter

A rain gutter is to be constructed of aluminum sheets 12 inches wide. After marking off a length of 4 inches from each edge, this length is bent up at an angle θ . See Figure 30. The area A of the opening may be expressed as a function of θ as

$$A(\theta) = 16\sin\theta(\cos\theta + 1)$$

Find the area A of the opening for $\theta = 30^\circ$, $\theta = 45^\circ$, and $\theta = 60^\circ$.

Solution For $\theta = 30^{\circ}$: $A(30^{\circ}) = 16 \sin 30^{\circ} (\cos 30^{\circ} + 1)$


$$= 16\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2} + 1\right) = 4\sqrt{3} + 8 \approx 14.9$$

The area of the opening for $\theta = 30^{\circ}$ is about 14.9 square inches.

-1

Figure 29

Figure 30



EXAMPLE 8

For
$$\theta = 45^{\circ}$$
: $A(45^{\circ}) = 16 \sin 45^{\circ} (\cos 45^{\circ} + 1)$
= $16 \left(\frac{\sqrt{2}}{2}\right) \left(\frac{\sqrt{2}}{2} + 1\right) = 8 + 8\sqrt{2} \approx 19.3$

The area of the opening for $\theta = 45^{\circ}$ is about 19.3 square inches.

For $\theta = 60^{\circ}$: $A(60^{\circ}) = 16 \sin 60^{\circ} (\cos 60^{\circ} + 1)$

$$= 16\left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{2}+1\right) = 12\sqrt{3} \approx 20.8$$

The area of the opening for $\theta = 60^{\circ}$ is about 20.8 square inches.

5 Find the Exact Values of the Trigonometric Functions for Integer Multiples of $\frac{\pi}{6} = 30^{\circ}, \frac{\pi}{4} = 45^{\circ}, \text{ and } \frac{\pi}{3} = 60^{\circ}$

We know the exact values of the trigonometric functions of $\frac{\pi}{4} = 45^{\circ}$. Using symmetry, we can find the exact values of the trigonometric functions of $\frac{3\pi}{4} = 135^{\circ}, \frac{5\pi}{4} = 225^{\circ}, \text{ and } \frac{7\pi}{4} = 315^{\circ}$. Figure 31 shows how.

As Figure 31 shows, using symmetry with respect to the *y*-axis, the point $\left(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $\frac{3\pi}{4} = 135^{\circ}$. Similarly, using symmetry with respect to the origin, the point $\left(-\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $\frac{5\pi}{4} = 225^{\circ}$. Finally, using symmetry with respect to the *x*-axis, the point $\left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $\frac{5\pi}{4} = 225^{\circ}$. Finally, using symmetry with respect to the *x*-axis, the point $\left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $\frac{7\pi}{4} = 315^{\circ}$.


EXAMPLE 9

Finding Exact Values for Multiples of $\frac{\pi}{4} = 45^{\circ}$

Based on Figure 31, we see that


(a)
$$\sin 135^\circ = \frac{\sqrt{2}}{2}$$
 (b) $\cos \frac{5\pi}{4} = -\frac{\sqrt{2}}{2}$ (c) $\tan 315^\circ = \frac{-\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = -1$

Figure 31 can also be used to find exact values for other multiples of $\frac{\pi}{4} = 45^{\circ}$. For example, the point $\left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $-\frac{\pi}{4} = -45^{\circ}$; the point $\left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ is the point on the unit circle that corresponds to the angle $\frac{9\pi}{4} = 405^{\circ}$.





The use of symmetry also provides information about certain integer multiples of the angles $\frac{\pi}{6} = 30^{\circ}$ and $\frac{\pi}{3} = 60^{\circ}$. See Figures 32 and 33.



EXAMPLE 10 Using Figures 32 and 33

Based on Figures 32 and 33, we see that

(a)
$$\cos 210^\circ = -\frac{\sqrt{3}}{2}$$
 (b) $\sin(-60^\circ) = -\frac{\sqrt{3}}{2}$ (c) $\tan \frac{5\pi}{3} = \frac{-\frac{\sqrt{3}}{2}}{\frac{1}{2}} = -\sqrt{3}$

6 Use a Calculator to Approximate the Value of a Trigonometric Function

NOW WORK PROBLEM 49.

Before getting started, you must first decide whether to enter the angle in the calculator using radians or degrees and then set the calculator to the correct MODE. Check your instruction manual to find out how your calculator handles degrees and radians. Your calculator has keys marked \sin , \cos , and \tan . To find the values of the remaining three trigonometric functions, secant, cosecant, and cotangent, we use the fact that, if P = (x, y) is a point on the unit circle on the terminal side of θ , then

$$\sec \theta = \frac{1}{x} = \frac{1}{\cos \theta}$$
 $\csc \theta = \frac{1}{y} = \frac{1}{\sin \theta}$ $\cot \theta = \frac{x}{y} = \frac{1}{\frac{y}{x}} = \frac{1}{\tan \theta}$

(c) $\tan \frac{\pi}{12}$

EXAMPLE 11

Using a Calculator to Approximate the Value of a Trigonometric Function

Use a calculator to find the approximate value of:

(a)
$$\cos 48^{\circ}$$
 (b) $\csc 21^{\circ}$

Express your answers rounded to two decimal places.

Solution (a) First we set the MODE to receive degrees. See Figure 34(a). Figure 34(b) shows the solution using a TI-84 Plus graphing calculator. Rounded to two decimal places,

 $\cos 48^\circ = 0.66991306 \approx 0.67$

Figure 34







Figure 35











- (b) Most calculators do not have a csc key. The manufacturers assume that the user knows some trigonometry. To find the value of $\csc 21^\circ$, use the fact that $\csc 21^\circ = \frac{1}{\sin 21^\circ}$. Figure 35 shows the solution using a TI-84 Plus graphing calculator. Rounded to two decimal places, $\csc 21^{\circ} \approx 2.79$
- (c) Set the MODE to receive radians. Figure 36 shows the solution using a TI-84 Plus graphing calculator. Rounded to two decimal places,

$$\operatorname{an}\frac{\pi}{12} \approx 0.27$$

Use a Circle of Radius r to Evaluate

NOW WORK PROBLEM 67.

the Trigonometric Functions

Until now, to find the exact value of a trigonometric function of an angle θ required that we locate the corresponding point P = (x, y) on the unit circle. In fact, though, any circle whose center is at the origin can be used.

Let θ be any nonquadrantal angle placed in standard position. Let P = (x, y) be the point on the circle $x^2 + y^2 = r^2$ that corresponds to θ , and let $P^* = (x^*, y^*)$ be the point on the unit circle that corresponds to θ . See Figure 37.

Notice that the triangles OA^*P^* and OAP are similar; as a result, the ratios of corresponding sides are equal.

$\frac{y^*}{1}$:	<u>y</u>	$\frac{x^{*}}{1} =$	<u>x</u>	$\frac{y^*}{y} = \frac{y}{y}$
1	r	1	r	$x^* x$
$\frac{1}{y^*}$	= r	$\frac{1}{x^*} =$	_ <u>r</u>	$\frac{x^*}{y^*} = \frac{x}{y}$
y^*	у	<i>x</i> *	х	<i>y</i> * <i>y</i>

These results lead us to formulate the following theorem:

Theorem

For an angle θ in standard position, let P = (x, y) be the point on the terminal side of θ that is also on the circle $x^2 + y^2 = r^2$. Then


$\sin\theta = \frac{y}{r}$	$\cos\theta = \frac{x}{r}$	$\tan \theta = \frac{y}{x}, x \neq 0$
$\csc \theta = \frac{r}{y}, y \neq 0$	$\sec \theta = \frac{r}{x}, x \neq 0$	$\cot \theta = \frac{x}{y}, y \neq 0$

EXAMPLE 12 Finding the Exact Values of the Six Trigonometric Functions

Find the exact values of each of the six trigonometric functions of an angle θ if (4, -3) is a point on its terminal side.

Solution See Figure 38. The point (4, -3) is on a circle of radius $r = \sqrt{4^2 + (-3)^2} = \sqrt{16 + 9} = \sqrt{25} = 5$ with the center at the origin.

Figure 38



For the point (x, y) = (4, -3), we have x = 4 and y = -3. Since r = 5, we find

$$\sin \theta = \frac{y}{r} = -\frac{3}{5} \qquad \cos \theta = \frac{x}{r} = \frac{4}{5} \qquad \tan \theta = \frac{y}{x} = -\frac{3}{4}$$
$$\csc \theta = \frac{r}{y} = -\frac{5}{3} \qquad \sec \theta = \frac{r}{x} = \frac{5}{4} \qquad \cot \theta = \frac{x}{y} = -\frac{4}{3}$$

HISTORICAL FEATURE


The name sine for the sine function is due to a medieval confusion. The name comes from the Sanskrit word *jiva* (meaning chord), first used in India by Araybhata the Elder (AD 510). He really meant halfchord, but abbreviated it. This was brought into Arabic as *jiba*, which was meaningless. Because the proper Arabic word *jaib* would be written the same way (short vowels are not written out in Arabic), *jiba* was pronounced as *jaib*, which meant bosom or hollow, and *jiba* remains as the Arabic word for sine to this day. Scholars translating the Arabic works into Latin found that the word *sinus* also meant bosom or hollow, and from *sinus* we get the word *sine*.

The name tangent, due to Thomas Finck (1583), can be understood by looking at Figure 39. The line segment \overline{DC} is tangent to the circle at C. If d(O, B) = d(O, C) = 1, then the length of the line segment \overline{DC} is

$$d(D, C) = \frac{d(D, C)}{1} = \frac{d(D, C)}{d(0, C)} = \tan \alpha$$

The old name for the tangent is *umbra versa* (meaning turned shadow), referring to the use of the tangent in solving height problems with shadows.

The names of the remaining functions came about as follows. If α and β are complementary angles, then $\cos \alpha = \sin \beta$. Because β is the complement of α , it was natural to write the cosine of α as sin co α . Probably for reasons involving ease of pronunciation, the co migrated to the front, and then cosine received a three-letter abbreviation to match sin, sec, and tan. The two other cofunctions were similarly treated, except that the long forms cotan and cosec survive to this day in some countries.



5.2 Assess Your Understanding

'Are You Prepared?'

Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

- **1.** In a right triangle, with legs *a* and *b* and hypotenuse *c*, the Pythagorean Theorem states that _____. (p. 962)
- 2. The value of the function f(x) = 3x 7 at 5 is _____. (pp. 61–63)
- **3.** *True or False:* For a function y = f(x), for each x in the domain, there is exactly one element y in the range. (pp. 55–61)
- 4. What is the equation of the unit circle? (p. 45)

Concepts and Vocabulary

- 7. $\tan \frac{\pi}{4} + \sin 30^\circ =$ _____.
- 8. Using a calculator, sin 2 = _____, rounded to two decimal places.

- 5. What point is symmetric with respect to the y-axis to the point $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$? (pp. 17–19)
- 6. If (x, y) is a point on the unit circle in quadrant IV and if $x = \frac{\sqrt{3}}{2}$, what is y? (p. 45)
- **9.** *True or False:* Exact values can be found for the trigonometric functions of 60°.
- **10.** *True or False:* Exact values can be found for the sine of any angle.

Skill Building

.

In Problems 11–18, t is a real number and P = (x, y) is the point on the unit circle that corresponds to t. Find the exact values of the six trigonometric functions of t.

11.
$$\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$$

12. $\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)$
13. $\left(-\frac{2}{5}, \frac{\sqrt{21}}{5}\right)$
14. $\left(-\frac{1}{5}, \frac{2\sqrt{6}}{5}\right)$
15. $\left(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$
16. $\left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$
17. $\left(\frac{2\sqrt{2}}{3}, -\frac{1}{3}\right)$
18. $\left(-\frac{\sqrt{5}}{3}, -\frac{2}{3}\right)$

In Problems 19–28, find the exact value. Do not use a calculator.

19. $\sin \frac{11\pi}{2}$ 20. $\cos(7\pi)$ 21. $\tan(6\pi)$ 22. $\cot \frac{7\pi}{2}$ 23. $\csc \frac{11\pi}{2}$ 24. $\sec(8\pi)$ 25. $\cos\left(-\frac{3\pi}{2}\right)$ 26. $\sin(-3\pi)$ 27. $\sec(-\pi)$ 28. $\tan(-3\pi)$

In Problems 29–48, find the exact value of each expression. Do not use a calculator.

31. $\sin 90^{\circ} + \tan 45^{\circ}$ **29.** $\sin 45^\circ + \cos 60^\circ$ **30.** $\sin 30^{\circ} - \cos 45^{\circ}$ **32.** $\cos 180^\circ - \sin 180^\circ$ **33.** sin 45° cos 45° **35.** csc 45° tan 60° **36.** sec 30° cot 45° **34.** $\tan 45^{\circ} \cos 30^{\circ}$ **39.** $2\sin\frac{\pi}{3} - 3\tan\frac{\pi}{6}$ **40.** $2\sin\frac{\pi}{4} + 3\tan\frac{\pi}{4}$ **38.** $5 \cos 90^\circ - 8 \sin 270^\circ$ **37.** $4 \sin 90^\circ - 3 \tan 180^\circ$ **42.** $\tan \frac{\pi}{3} + \cos \frac{\pi}{3}$ **43.** $2 \sec \frac{\pi}{4} + 4 \cot \frac{\pi}{3}$ **44.** $3 \csc \frac{\pi}{3} + \cot \frac{\pi}{4}$ 41. $\sin \frac{\pi}{4} - \cos \frac{\pi}{4}$ **46.** $\sin \frac{3\pi}{2} + \tan \pi$ **47.** $\csc \frac{\pi}{2} + \cot \frac{\pi}{2}$ **48.** sec $\pi - \csc \frac{\pi}{2}$ **45.** $\tan \pi - \cos 0$

In Problems 49–66, find the exact values of the six trigonometric functions of the given angle. If any are not defined, say "not defined." Do not use a calculator.

49. $\frac{2\pi}{3}$	50. $\frac{5\pi}{6}$	51. 210°	52. 240°	53. $\frac{3\pi}{4}$	54. $\frac{11\pi}{4}$
55. $\frac{8\pi}{3}$	56. $\frac{13\pi}{6}$	57. 405°	58. 390°	59. $-\frac{\pi}{6}$	60. $-\frac{\pi}{3}$
61. -45°	62. −60°	63. $\frac{5\pi}{2}$	64. 5π	65. 720°	66. 630°

111. $f(-\theta)$ **112.** $g(-\theta)$

In Problems 67–82, use a calculator to find the approximate value of each expression rounded to two decimal places.					
67. sin 28°	68. cos 14°		69. tan 21°	70. cot 70°	
71. sec 41°	72. csc 55°		73. $\sin \frac{\pi}{10}$	74. $\cos \frac{\pi}{8}$	
75. $\tan \frac{5\pi}{12}$	76. $\cot \frac{\pi}{18}$		77. $\sec \frac{\pi}{12}$	78. $\csc \frac{5\pi}{13}$	
79. sin 1	80. tan 1	:	81. sin 1°	82. tan 1°	
In Problems 83–92,	a point on the terminal si	de of an angle θ is giver	n. Find the exact values	of the six trigonometric functions of θ .	
83. (-3, 4)	84. (5, -12)	85. (2, -3)	86. (-1, -2)	87. (-2, -2)	
88. (1, -1)	89. (-3, -2)	90. (2, 2)	91. $\left(\frac{1}{3}, -\frac{1}{4}\right)$	92. (-0.3, -0.4)	
93. Find the exact v	value of:	94	Find the exact value of	f:	
sin 45° ⊣	$-\sin 135^\circ + \sin 225^\circ + \sin 225^\circ$	n 315°	tan 6	50° + tan 150°	
95. If $\sin \theta = 0.1$, fi	$ind \sin(\theta + \pi).$	96	If $\cos \theta = 0.3$, find $\cos \theta$	$s(\theta + \pi).$	
97. If $\tan \theta = 3$, fin	$d\tan(\theta + \pi).$	98.	If $\cot \theta = -2$, find $\cot \theta$	$(\theta + \pi).$	
99. If $\sin \theta = \frac{1}{5}$, fin	d csc θ .	100.	If $\cos \theta = \frac{2}{3}$, find sec θ	9.	
In Problems 101–112	$g, f(\theta) = \sin \theta \text{ and } g(\theta) =$	$\cos \theta$. Find the exact val	ue of each function belo	w if $\theta = 60^{\circ}$. Do not use a calculator.	
101. <i>f</i> (<i>θ</i>)	102. $g(\theta)$ 103	$5. \ f\left(\frac{\theta}{2}\right) \qquad \qquad 104.$	$g\left(\frac{\theta}{2}\right)$ 105. [$f(\theta)]^2$ 106. $[g(\theta)]^2$	

 $\not \bigtriangleup 113$. Use a calculator in radian mode to complete the following table.

108. g(2 θ)

107. *f*(2*θ*)

What can you conclude about the value of $\frac{\sin \theta}{\theta}$ as θ approaches 0?

109. $2f(\theta)$


θ	0.5	0.4	0.2	0.1	0.01	0.001	0.0001	0.00001
$\sin \theta$								
$\frac{\sin \theta}{\theta}$								
θ								

110. $2g(\theta)$

What can you conclude about the value of $\frac{\cos \theta - 1}{\theta}$ as θ approaches 0?

θ	0.5	0.4	0.2	0.1	0.01	0.001	0.0001	0.00001
$\cos \theta - 1$								
$\frac{\cos\theta - 1}{\theta}$								

Projectile Motion The path of a projectile fired at an inclination θ to the horizontal with initial speed v_0 is a parabola (see the figure).



The range R of the projectile, that is, the horizontal distance that the projectile travels, is found by using the formula

$$R = \frac{v_0^2 \sin(2\theta)}{g}$$

where $g \approx 32.2$ feet per second per second ≈ 9.8 meters per second per second is the acceleration due to gravity. The maximum height H of the projectile is


$$H = \frac{v_0^2 (\sin \theta)^2}{2g}$$

In Problems 115–118, find the range R and maximum height H.

- **115.** The projectile is fired at an angle of 45° to the horizontal with an initial speed of 100 feet per second.
- **116.** The projectile is fired at an angle of 30° to the horizontal with an initial speed of 150 meters per second.
- **117.** The projectile is fired at an angle of 25° to the horizontal with an initial speed of 500 meters per second.
- **118.** The projectile is fired at an angle of 50° to the horizontal with an initial speed of 200 feet per second.
- **119. Inclined Plane** If friction is ignored, the time *t* (in seconds) required for a block to slide down an inclined plane (see the figure) is given by the formula

$$t = \sqrt{\frac{2a}{g\sin\theta\cos\theta}}$$


where *a* is the length (in feet) of the base and $g \approx 32$ feet per second per second is the acceleration due to gravity. How long does it take a block to slide down an inclined plane with base a = 10 feet when:



120. Piston Engines In a certain piston engine, the distance x (in centimeters) from the center of the drive shaft to the head of the piston is given by

$$x = \cos\theta + \sqrt{16 + 0.5\cos(2\theta)}$$


where θ is the angle between the crank and the path of the piston head (see the figure). Find *x* when $\theta = 30^{\circ}$ and when $\theta = 45^{\circ}$.



121. Calculating the Time of a Trip Two oceanfront homes are located 8 miles apart on a straight stretch of beach, each a distance of 1 mile from a paved road that parallels the ocean. Sally can jog 8 miles per hour along the paved road, but only 3 miles per hour in the sand on the beach. Because of a river directly between the two houses, it is necessary to jog in the sand to the road, continue on the road, and then jog directly back in the sand to get from one house to the other. See the illustration. The time *T* to get from one house to the other as a function of the angle θ shown in the illustration is

$$T(\theta) = 1 + \frac{2}{3\sin\theta} - \frac{1}{4\tan\theta}, \qquad 0^{\circ} < \theta < 90^{\circ}$$

- (a) Calculate the time T for $\theta = 30^{\circ}$. How long is Sally on the paved road?
- (b) Calculate the time T for $\theta = 45^{\circ}$. How long is Sally on the paved road?
- (c) Calculate the time T for $\theta = 60^{\circ}$. How long is Sally on the paved road?
- (d) Calculate the time T for $\theta = 90^{\circ}$. Describe the path taken. Why can't the formula for T be used?




122. Designing Fine Decorative Pieces A designer of decorative art plans to market solid gold spheres encased in clear crystal cones. Each sphere is of fixed radius R and will be enclosed in a cone of height h and radius r. See the illustration. Many cones can be used to enclose the sphere, each having a different slant angle θ . The volume V of the cone can be expressed as a function of the slant angle θ of the cone as

SECTION 5.2 Trigonometric Functions: Unit Circle Approach 387

$$V(\theta) = \frac{1}{3}\pi R^3 \frac{(1 + \sec \theta)^3}{(\tan \theta)^2}, \qquad 0^\circ < \theta < 90^\circ$$


What volume V is required to enclose a sphere of radius 2 centimeters in a cone whose slant angle θ is 30°? 45°? 60°?



123. Projectile Motion An object is propelled upward at an angle θ , $45^{\circ} < \theta < 90^{\circ}$, to the horizontal with an initial velocity of v_0 feet per second from the base of an inclined plane that makes an angle of 45° with the horizontal. See the illustration. If air resistance is ignored, the distance *R* that it travels up the inclined plane is given by

$$R = \frac{v_0^2 \sqrt{2}}{32} [\sin(2\theta) - \cos(2\theta) - 1]$$

- (a) Find the distance *R* that the object travels along the inclined plane if the initial velocity is 32 feet per second and $\theta = 60^{\circ}$.
- (b) Graph $R = R(\theta)$ if the initial velocity is 32 feet per second.
- (c) What value of θ makes *R* largest?



Discussion and Writing


- **129.** Write a brief paragraph that explains how to quickly compute the trigonometric functions of 30° , 45° , and 60° .
- **130.** Write a brief paragraph that explains how to quickly compute the trigonometric functions of 0°, 90°, 180°, and 270°.

'Are You Prepared?' Answers


1.
$$c^2 = a^2 + b^2$$
 2. 8 **3.** True **4.** $x^2 + y^2 = 1$ **5.** $\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ **6.** $-\frac{1}{2}$

124. If $\theta, 0 < \theta < \pi$, is the angle between the positive *x*-axis and a nonhorizontal, nonvertical line *L*, show that the slope *m* of *L* equals tan θ . The angle θ is called the **inclination** of *L*.

[Hint: See the illustration, where we have drawn the line M parallel to L and passing through the origin. Use the fact that M intersects the unit circle at the point $(\cos \theta, \sin \theta)$.]



In Problems 125–128, use the figure to approximate the value of the six trigonometric functions at t to the nearest tenth. Then use a calculator to approximate each of the six trigonometric functions at t.



131. How would you explain the meaning of the sine function to a fellow student who has just completed college algebra?

5.3 Properties of the Trigonometric Functions

PREPARING FOR THIS SECTION Before getting started, review the following:

- Functions (Section 2.1, pp. 56-65)
- Identity (Appendix, Section A.5, p. 984)

Now work the 'Are You Prepared?' problems on page 399.

OBJECTIVES 1 Determine the Domain and the Range of the Trigonometric Functions

- 2 Determine the Period of the Trigonometric Functions
- ³ Determine the Signs of the Trigonometric Functions in a Given Quadrant
- 4 Find the Values of the Trigonometric Functions Using Fundamental Identities
- 5 Find the Exact Values of the Trigonometric Functions of an Angle Given One of the Functions and the Quadrant of the Angle

• Even and Odd Functions (Section 2.3, pp. 80–82)

6 Use Even–Odd Properties to Find the Exact Values of the Trigonometric Functions

J Determine the Domain and the Range of the Trigonometric Functions

Figure 40 P = (x, y) (0, 1) (-1, 0) (0, -1) (1, 0) (1, 0) (0, -1) Let θ be an angle in standard position, and let P = (x, y) be the point on the unit circle that corresponds to θ . See Figure 40. Then, by definition,

$$\sin \theta = y \qquad \cos \theta = x \qquad \tan \theta = \frac{y}{x}, \quad x \neq 0$$
$$\csc \theta = \frac{1}{y}, \quad y \neq 0 \qquad \sec \theta = \frac{1}{x}, \quad x \neq 0 \qquad \cot \theta = \frac{x}{y}, \quad y \neq 0$$

For sin θ and cos θ , θ can be any angle, so it follows that the domain of the sine function and cosine function is the set of all real numbers.

The domain of the sine function is the set of all real numbers. The domain of the cosine function is the set of all real numbers.

If x = 0, then the tangent function and the secant function are not defined. That is, for the tangent function and secant function, the *x*-coordinate of P = (x, y) cannot be 0. On the unit circle, there are two such points (0, 1) and (0, -1). These two points correspond to the angles $\frac{\pi}{2}(90^\circ)$ and $\frac{3\pi}{2}(270^\circ)$ or, more generally, to any angle that is an odd integer multiple of $\frac{\pi}{2}(90^\circ)$, such as $\frac{\pi}{2}(90^\circ)$, $\frac{3\pi}{2}(270^\circ)$, $\frac{5\pi}{2}(450^\circ)$, $-\frac{\pi}{2}(-90^\circ)$, $-\frac{3\pi}{2}(-270^\circ)$, and so on. Such angles must therefore be excluded from the domain of the tangent function and secant function. The domain of the tangent function is the set of all real numbers, except odd integer multiples of $\frac{\pi}{2}(90^{\circ})$.

The domain of the secant function is the set of all real numbers, except odd integer multiples of $\frac{\pi}{2}(90^{\circ})$.

If y = 0, then the cotangent function and the cosecant function are not defined. For the cotangent function and cosecant function, the *y*-coordinate of P = (x, y) cannot be 0. On the unit circle, there are two such points, (1, 0) and (-1, 0). These two points correspond to the angles $0(0^{\circ})$ and $\pi(180^{\circ})$ or, more generally, to any angle that is an integer multiple of $\pi(180^{\circ})$, such as $0(0^{\circ})$, $\pi(180^{\circ})$, $2\pi(360^{\circ})$, $3\pi(540^{\circ})$, $-\pi(-180^{\circ})$, and so on. Such angles must therefore be excluded from the domain of the cotangent function and cosecant function.

The domain of the cotangent function is the set of all real numbers, except integer multiples of $\pi(180^\circ)$.

The domain of the cosecant function is the set of all real numbers, except integer multiples of $\pi(180^\circ)$.

Next we determine the range of each of the six trigonometric functions. Refer again to Figure 40. Let P = (x, y) be the point on the unit circle that corresponds to the angle θ . It follows that $-1 \le x \le 1$ and $-1 \le y \le 1$. Consequently, since $\sin \theta = y$ and $\cos \theta = x$, we have

 $-1 \le \sin \theta \le 1$ $-1 \le \cos \theta \le 1$

The range of both the sine function and the cosine function consists of all real numbers between -1 and 1, inclusive. Using absolute value notation, we have $|\sin \theta| \le 1$ and $|\cos \theta| \le 1$.

If θ is not an integer multiple of $\pi(180^\circ)$, then $\csc \theta = \frac{1}{y}$. Since $y = \sin \theta$ and $|y| = |\sin \theta| \le 1$, it follows that $|\csc \theta| = \frac{1}{|\sin \theta|} = \frac{1}{|y|} \ge 1$. The range of the cosecant function consists of all real numbers less than or equal to -1 or greater than or equal to 1. That is,

 $\csc \theta \leq -1$ or $\csc \theta \geq 1$

If θ is not an odd integer multiple of $\frac{\pi}{2}(90^\circ)$, then, by definition, sec $\theta = \frac{1}{x}$. Since $x = \cos \theta$ and $|x| = |\cos \theta| \le 1$, it follows that $|\sec \theta| = \frac{1}{|\cos \theta|} = \frac{1}{|x|} \ge 1$. The range of the secant function consists of all real numbers less than or equal to -1 or greater than or equal to 1. That is,

$$\sec \theta \leq -1$$
 or $\sec \theta \geq 1$

The range of both the tangent function and the cotangent function is the set of all real numbers. You are asked to prove this in Problems 121 and 122.

 $-\infty < \tan \theta < \infty$ $-\infty < \cot \theta < \infty$

Table 4 summarizes these results.


Table 4

Function	Symbol	Domain	Range
sine	$f(\theta) = \sin \theta$	All real numbers	All real numbers from -1 to 1, inclusive
cosine	$f(\theta) = \cos \theta$	All real numbers	All real numbers from -1 to 1, inclusive
tangent	f(heta) = an heta	All real numbers, except odd integer multiples of $rac{\pi}{2}(90^\circ)$	All real numbers
cosecant	$f(\theta) = \csc \theta$	All real numbers, except integer multiples of $\pi(180^\circ)$	All real numbers greater than or equal to 1 or less than or equal to -1
secant	$f(heta) = \sec heta$	All real numbers, except odd integer multiples of $\frac{\pi}{2}$ (90°)	All real numbers greater than or equal to 1 or less than or equal to -1
cotangent	$f(\theta) = \cot \theta$	All real numbers, except integer multiples of $\pi(180^\circ)$	All real numbers

NOW WORK PROBLEM 97.

Determine the Period of the Trigonometric Functions


Figure 41



Look at Figure 41. This figure shows that for an angle of $\frac{\pi}{3}$ radians the corresponding point *P* on the unit circle is $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. Notice that, for an angle of $\frac{\pi}{3} + 2\pi$ radians, the corresponding point *P* on the unit circle is also $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. Then

$$\sin\frac{\pi}{3} = \frac{\sqrt{3}}{2} \quad \text{and} \quad \sin\left(\frac{\pi}{3} + 2\pi\right) = \frac{\sqrt{3}}{2}$$
$$\cos\frac{\pi}{3} = \frac{1}{2} \quad \text{and} \quad \cos\left(\frac{\pi}{3} + 2\pi\right) = \frac{1}{2}$$

Figure 42



This example illustrates a more general situation. For a given angle θ , measured in radians, suppose that we know the corresponding point P = (x, y) on the unit circle. Now add 2π to θ . The point on the unit circle corresponding to $\theta + 2\pi$ is identical to the point P on the unit circle corresponding to θ . See Figure 42. The values of the trigonometric functions of $\theta + 2\pi$ are equal to the values of the corresponding trigonometric functions of θ .

If we add (or subtract) integer multiples of 2π to θ , the trigonometric values remain unchanged. That is, for all θ .

 $\sin(\theta + 2\pi k) = \sin \theta \qquad \cos(\theta + 2\pi k) = \cos \theta$ where k is any integer. (1)

Functions that exhibit this kind of behavior are called *periodic functions*.

A function f is called **periodic** if there is a positive number p such that, whenever θ is in the domain of f, so is $\theta + p$, and

$$f(\theta + p) = f(\theta)$$

If there is a smallest such number p, this smallest value is called the **(fundamental) period** of f.

Based on equation (1), the sine and cosine functions are periodic. In fact, the sine and cosine functions have period 2π . You are asked to prove this fact in Problems 123 and 124. The secant and cosecant functions are also periodic with period 2π , and the tangent and cotangent functions are periodic with period π . You are asked to prove these statements in Problems 125 through 128.

These facts are summarized as follows:

Periodic Properties

 $\sin(\theta + 2\pi) = \sin\theta \quad \cos(\theta + 2\pi) = \cos\theta \quad \tan(\theta + \pi) = \tan\theta$ $\csc(\theta + 2\pi) = \csc\theta \quad \sec(\theta + 2\pi) = \sec\theta \quad \cot(\theta + \pi) = \cot\theta$

Because the sine, cosine, secant, and cosecant functions have period 2π , once we know their values for $0 \le \theta < 2\pi$, we know all their values; similarly, since the tangent and cotangent functions have period π , once we know their values for $0 \le \theta < \pi$, we know all their values.

EXAMPLE 1 Finding Exact Values Using Periodic Properties


Find the exact value of:

(a)
$$\sin \frac{17\pi}{4}$$
 (b) $\cos(5\pi)$ (c) $\tan \frac{5\pi}{4}$

Solution

(a) It is best to sketch the angle first, as shown in Figure 43(a). Since the period of the sine function is 2π , each full revolution can be ignored. This leaves the angle $\frac{\pi}{4}$. Then

$$\sin\frac{17\pi}{4} = \sin\left(\frac{\pi}{4} + 4\pi\right) = \sin\frac{\pi}{4} = \frac{\sqrt{2}}{2}$$



In Words Tangent and cotangent have

period π ; the others have period 2π .

(b) See Figure 43(b). Since the period of the cosine function is 2π , each full revolution can be ignored. This leaves the angle π . Then

$$\cos(5\pi) = \cos(\pi + 4\pi) = \cos\pi = -1$$

(c) See Figure 43(c). Since the period of the tangent function is π , each half-

revolution can be ignored. This leaves the angle $\frac{\pi}{4}$. Then

$$\tan\frac{5\pi}{4} = \tan\left(\frac{\pi}{4} + \pi\right) = \tan\frac{\pi}{4} = 1$$

11

The periodic properties of the trigonometric functions will be very helpful to us when we study their graphs later in the chapter.

NOW WORK PROBLEM 11.


Jost Determine the Signs of the Trigonometric Functions in a Given Quadrant

Let P = (x, y) be the point on the unit circle that corresponds to the angle θ . If we know in which quadrant the point P lies, then we can determine the signs of the trigonometric functions of θ . For example, if P = (x, y) lies in quadrant IV, as shown in Figure 44, then we know that x > 0 and y < 0. Consequently,


$$\sin \theta = y < 0 \qquad \cos \theta = x > 0 \qquad \tan \theta = \frac{y}{x} < 0$$
$$\csc \theta = \frac{1}{y} < 0 \qquad \sec \theta = \frac{1}{x} > 0 \qquad \cot \theta = \frac{x}{y} < 0$$

Table 5 lists the signs of the six trigonometric functions for each quadrant. See also Figure 45.

Table 5	Quadrant of P	$\sin \theta$, csc θ	$\cos \theta$, $\sec \theta$	$\tan \theta$, $\cot \theta$
	I	Positive	Positive	Positive
	II	Positive	Negative	Negative
	III	Negative	Negative	Positive
	IV	Negative	Positive	Negative







EXAMPLE 2 Finding the Quadrant in Which an Angle θ Lies

If $\sin \theta < 0$ and $\cos \theta < 0$, name the quadrant in which the angle θ lies.

Solution Let P = (x, y) be the point on the unit circle corresponding to θ . Then $\sin \theta = y < 0$ and $\cos \theta = x < 0$. The point P = (x, y) must be in quadrant III, so θ lies in quadrant III.

NOW WORK PROBLEM 27.

Find the Values of the Trigonometric Functions Using Fundamental Identities

If P = (x, y) is the point on the unit circle corresponding to θ , then

$$\sin \theta = y \qquad \cos \theta = x \qquad \tan \theta = \frac{y}{x}, \quad \text{if } x \neq 0$$
$$\csc \theta = \frac{1}{y}, \quad \text{if } y \neq 0 \qquad \sec \theta = \frac{1}{x}, \quad \text{if } x \neq 0 \qquad \cot \theta = \frac{x}{y}, \quad \text{if } y \neq 0$$

Based on these definitions, we have the **reciprocal identities:**

Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \cot \theta = \frac{1}{\tan \theta}$$
 (2)

Two other fundamental identities are the quotient identities.

Quotient Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$
(3)

The proofs of identities (2) and (3) follow from the definitions of the trigonometric functions. (See Problems 129 and 130.)

If $\sin \theta$ and $\cos \theta$ are known, identities (2) and (3) make it easy to find the values of the remaining trigonometric functions.

EXAMPLE 3 Finding Exact Values Using Identities When Sine and Cosine Are Given

Given $\sin \theta = \frac{\sqrt{5}}{5}$ and $\cos \theta = \frac{2\sqrt{5}}{5}$, find the exact values of the four remaining trigonometric functions of θ using identities.

Solution Based on a quotient identity from (3), we have

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{\sqrt{5}}{5}}{\frac{2\sqrt{5}}{5}} = \frac{1}{2}$$

Then we use the reciprocal identities from (2) to get

$$\csc \theta = \frac{1}{\sin \theta} = \frac{1}{\frac{\sqrt{5}}{5}} = \frac{5}{\sqrt{5}} = \sqrt{5} \qquad \sec \theta = \frac{1}{\cos \theta} = \frac{1}{\frac{2\sqrt{5}}{5}} = \frac{5}{2\sqrt{5}} = \frac{\sqrt{5}}{2} \qquad \cot \theta = \frac{1}{\tan \theta} = \frac{1}{\frac{1}{2}} = 2$$

NOW WORK PROBLEM 35.

The equation of the unit circle is $x^2 + y^2 = 1$, or equivalently, $y^2 + x^2 = 1$. If P = (x, y) is the point on the unit circle that corresponds to the angle θ , then

$$y^2 + x^2 = 1$$

But $y = \sin \theta$ and $x = \cos \theta$, so

$$(\sin\theta)^2 + (\cos\theta)^2 = 1 \tag{4}$$

It is customary to write $\sin^2 \theta$ instead of $(\sin \theta)^2$, $\cos^2 \theta$ instead of $(\cos \theta)^2$, and so on. With this notation, we can rewrite equation (4) as

$$\sin^2\theta + \cos^2\theta = 1$$
 (5)

If $\cos \theta \neq 0$, we can divide each side of equation (5) by $\cos^2 \theta$.

$$\frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} = \frac{1}{\cos^2 \theta}$$
$$\left(\frac{\sin \theta}{\cos \theta}\right)^2 + 1 = \left(\frac{1}{\cos \theta}\right)^2$$

Now use identities (2) and (3) to get

$$\tan^2\theta + 1 = \sec^2\theta \tag{6}$$

Similarly, if $\sin \theta \neq 0$, we can divide equation (5) by $\sin^2 \theta$ and use identities (2) and (3) to get $1 + \cot^2 \theta = \csc^2 \theta$, which we write as

$$\cot^2\theta + 1 = \csc^2\theta \tag{7}$$

Collectively, the identities in (5), (6), and (7) are referred to as the **Pythagorean** identities.

Let's pause here to summarize the fundamental identities.

Fundamental Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$
$$\csc \theta = \frac{1}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \cot \theta = \frac{1}{\tan \theta}$$
$$\sin^2 \theta + \cos^2 \theta = 1 \qquad \tan^2 \theta + 1 = \sec^2 \theta \qquad \cot^2 \theta + 1 = \csc^2 \theta$$

EXAMPLE 4 Finding the Exact Value of a Trigonometric Expression Using Identities

Find the exact value of each expression. Do not use a calculator.

(a)
$$\tan 20^{\circ} - \frac{\sin 20^{\circ}}{\cos 20^{\circ}}$$
 (b) $\sin^2 \frac{\pi}{12} + \frac{1}{\sec^2 \frac{\pi}{12}}$
Solution (a) $\tan 20^{\circ} - \frac{\sin 20^{\circ}}{\cos 20^{\circ}} = \tan 20^{\circ} - \tan 20^{\circ} = 0$
 \uparrow
 $\frac{\sin \theta}{\cos \theta} = \tan \theta$
(b) $\sin^2 \frac{\pi}{12} + \frac{1}{\sec^2 \frac{\pi}{12}} = \sin^2 \frac{\pi}{12} + \cos^2 \frac{\pi}{12} = 1$
 $\cos \theta = \frac{1}{\sec \theta}$ $\sin^2 \theta + \cos^2 \theta = 1$

Find the Exact Values of the Trigonometric Functions of an Angle Given One of the Functions and the Quadrant of the Angle

Many problems require finding the exact values of the remaining trigonometric functions when the value of one of them is known and the quadrant in which θ lies can be found. There are two approaches to solving such problems. One approach uses a circle of radius *r*, the other uses identities.

When using identities, sometimes a rearrangement is required. For example, the Pythagorean identity

$$\sin^2\theta + \cos^2\theta = 1$$

can be solved for sin θ in terms of cos θ (or vice versa) as follows:

$$\sin^2 \theta = 1 - \cos^2 \theta$$
$$\sin \theta = \pm \sqrt{1 - \cos^2 \theta}$$

where the + sign is used if $\sin \theta > 0$ and the - sign is used if $\sin \theta < 0$. Similarly, in $\tan^2 \theta + 1 = \sec^2 \theta$, we can solve for $\tan \theta$ (or $\sec \theta$), and in $\cot^2 \theta + 1 = \csc^2 \theta$, we can solve for $\cot \theta$ (or $\csc \theta$).


EXAMPLE 5

Finding Exact Values Given One Value and the Sign of Another

Given that $\sin \theta = \frac{1}{3}$ and $\cos \theta < 0$, find the exact values of each of the remaining five trigonometric functions.

Solution 1 Using a Circle

Figure 46



Suppose that P = (x, y) is the point on a circle that corresponds to θ . Since $\sin \theta = \frac{1}{3} > 0$ and $\cos \theta < 0$, the point P = (x, y) is in quadrant II. Because $\sin \theta = \frac{1}{3} = \frac{y}{r}$, we let y = 1 and r = 3. The point P = (x, y) that corresponds to θ lies on the circle of radius 3, namely $x^2 + y^2 = 9$. See Figure 46.

To find x, we use the fact that $x^2 + y^2 = 9$, y = 1, and P is in quadrant II.

$$x^{2} + y^{2} = 9$$

$$x^{2} + 1^{2} = 9$$

$$x^{2} = 8$$

$$x = -2\sqrt{2}$$

$$x < 0$$

Since $x = -2\sqrt{2}$ y = 1, and r = 3, we find that

$$\cos \theta = \frac{x}{r} = -\frac{2\sqrt{2}}{3} \qquad \tan \theta = \frac{y}{x} = \frac{1}{-2\sqrt{2}} = -\frac{\sqrt{2}}{4}$$
$$\csc \theta = \frac{r}{y} = \frac{3}{1} = 3 \qquad \sec \theta = \frac{r}{x} = \frac{3}{-2\sqrt{2}} = -\frac{3\sqrt{2}}{4} \qquad \cot \theta = \frac{x}{y} = \frac{-2\sqrt{2}}{1} = -2\sqrt{2}$$

Solution 2 Using Identities

First, we solve equation (5) for
$$\cos \theta$$
.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\cos^2 \theta = 1 - \sin^2 \theta$$

 $\cos\theta = \pm\sqrt{1-\sin^2\theta}$

Because $\cos \theta < 0$, we choose the minus sign.

$$\cos \theta = -\sqrt{1 - \sin^2 \theta} = -\sqrt{1 - \frac{1}{9}} = -\sqrt{\frac{8}{9}} = -\frac{2\sqrt{2}}{3}$$
$$\sin \theta = \frac{1}{3}$$

Now we know the values of $\sin \theta$ and $\cos \theta$, so we can use identities (2) and (3) to get

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{1}{3}}{\frac{-2\sqrt{2}}{3}} = \frac{1}{-2\sqrt{2}} = -\frac{\sqrt{2}}{4} \qquad \cot \theta = \frac{1}{\tan \theta} = -2\sqrt{2}$$
$$\sec \theta = \frac{1}{\cos \theta} = \frac{1}{\frac{-2\sqrt{2}}{3}} = \frac{-3}{2\sqrt{2}} = -\frac{3\sqrt{2}}{4} \qquad \csc \theta = \frac{1}{\sin \theta} = \frac{1}{\frac{1}{3}} = 3$$

Finding the Values of the Trigonometric Functions When One Is Known

Given the value of one trigonometric function and the quadrant in which θ lies, the exact value of each of the remaining five trigonometric functions can be found in either of two ways.

Method 1 Using a Circle of Radius r

- **STEP 1:** Draw a circle showing the location of the angle θ and the point P = (x, y) that corresponds to θ . The radius of the circle is $r = \sqrt{x^2 + y^2}$.
- **STEP 2:** Assign a value to two of the three variables x, y, r based on the value of the given trigonometric function and the location of P.
- **STEP 3:** Use the fact that *P* lies on the circle $x^2 + y^2 = r^2$ to find the value of the missing variable.
- **STEP 4:** Apply the theorem on page 382 to find the values of the remaining trigonometric functions.

Method 2 Using Identities

Use appropriately selected identities to find the value of each of the remaining trigonometric functions.

EXAMPLE 6 Given One Value of a Trigonometric Function, Find the Remaining Ones

Given that $\tan \theta = \frac{1}{2}$ and $\sin \theta < 0$, find the exact value of each of the remaining five trigonometric functions of θ .

Solution 1 Using a Circle

STEP 1: Since $\tan \theta = \frac{1}{2} > 0$ and $\sin \theta < 0$, the point P = (x, y) that corresponds to θ lies in quadrant III. See Figure 47.


Figure 47

STEP 2: Since $\tan \theta = \frac{1}{2} = \frac{y}{x}$ and θ lies in quadrant III, we let x = -2 and y = -1.

STEP 3: Then $r = \sqrt{x^2 + y^2} = \sqrt{(-2)^2 + (-1)^2} = \sqrt{5}$, and *P* lies on the circle $x^2 + y^2 = 5$.

STEP 4: Now apply the theorem on p. 382 using x = -2, y = -1, $r = \sqrt{5}$.

$$\sin \theta = \frac{y}{r} = \frac{-1}{\sqrt{5}} = -\frac{\sqrt{5}}{5} \qquad \cos \theta = \frac{x}{r} = \frac{-2}{\sqrt{5}} = -\frac{2\sqrt{5}}{5}$$
$$\csc \theta = \frac{r}{y} = \frac{\sqrt{5}}{-1} = -\sqrt{5} \qquad \sec \theta = \frac{r}{x} = \frac{\sqrt{5}}{-2} = -\frac{\sqrt{5}}{2} \qquad \cot \theta = \frac{x}{y} = \frac{-2}{-1} = 2$$



Solution 2 Using Identities

We use the Pythagorean identity that involves $\tan \theta$, that is, $\tan^2 \theta + 1 = \sec^2 \theta$. Since $\tan \theta = \frac{1}{2} > 0$ and $\sin \theta < 0$, then θ lies in quadrant III, where $\sec \theta < 0$.

$$\tan^{2} \theta + 1 = \sec^{2} \theta$$
Pythagorean identity
$$\left(\frac{1}{2}\right)^{2} + 1 = \sec^{2} \theta$$

$$\tan \theta = \frac{1}{2}$$

$$\sec^{2} \theta = \frac{1}{4} + 1 = \frac{5}{4}$$
Proceed to solve for sec θ .
$$\sec \theta = -\frac{\sqrt{5}}{2}$$

$$\sec \theta < 0$$

Then

$$\cos \theta = \frac{1}{\sec \theta} = \frac{1}{-\frac{\sqrt{5}}{2}} = -\frac{2}{\sqrt{5}} = -\frac{2\sqrt{5}}{5}$$
$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \text{so} \quad \sin \theta = \tan \theta \cdot \cos \theta = \left(\frac{1}{2}\right) \cdot \left(-\frac{2\sqrt{5}}{5}\right) = -\frac{\sqrt{5}}{5}$$
$$\csc \theta = \frac{1}{\sin \theta} = \frac{1}{-\frac{\sqrt{5}}{5}} = -\frac{5}{\sqrt{5}} = -\sqrt{5}$$
$$\cot \theta = \frac{1}{\tan \theta} = \frac{1}{\frac{1}{2}} = 2$$

6 Use Even–Odd Properties to Find the Exact Values of the Trigonometric Functions

Recall that a function f is even if $f(-\theta) = f(\theta)$ for all θ in the domain of f; a function f is odd if $f(-\theta) = -f(\theta)$ for all θ in the domain of f. We will now show that the trigonometric functions sine, tangent, cotangent, and cosecant are odd functions and the functions cosine and secant are even functions.

Theorem

In Words

Cosine and secant are even functions; the others are odd functions.

Even-Odd Properties

 $\sin(-\theta) = -\sin\theta \qquad \cos(-\theta) = \cos\theta \qquad \tan(-\theta) = -\tan\theta$ $\csc(-\theta) = -\csc\theta \qquad \sec(-\theta) = \sec\theta \qquad \cot(-\theta) = -\cot\theta$

Proof Let P = (x, y) be the point on the unit circle that corresponds to the angle θ . See Figure 48. The point Q on the unit circle that corresponds to the angle $-\theta$ will have coordinates (x, -y). Using the definition of the trigonometric functions, we have

 $\sin \theta = y$ $\sin(-\theta) = -y$ $\cos \theta = x$ $\cos(-\theta) = x$

Figure 48

so

$$\sin(-\theta) = -y = -\sin\theta$$
 $\cos(-\theta) = x = \cos\theta$

Now, using these results and some of the fundamental identities, we have


$$\tan(-\theta) = \frac{\sin(-\theta)}{\cos(-\theta)} = \frac{-\sin\theta}{\cos\theta} = -\tan\theta \qquad \cot(-\theta) = \frac{1}{\tan(-\theta)} = \frac{1}{-\tan\theta} = -\cot\theta$$
$$\sec(-\theta) = \frac{1}{\cos(-\theta)} = \frac{1}{\cos\theta} = \sec\theta \qquad \csc(-\theta) = \frac{1}{\sin(-\theta)} = \frac{1}{-\sin\theta} = -\csc\theta$$

EXAMPLE 7 Finding Exact Values Using Even–Odd Properties

Find the exact value of:

(a)
$$\sin(-45^{\circ})$$
 (b) $\cos(-\pi)$ (c) $\cot\left(-\frac{3\pi}{2}\right)$ (d) $\tan\left(-\frac{37\pi}{4}\right)$

Solution



NOW WORK PROBLEM 59.

Odd function

5.3 Assess Your Understanding

'Are You Prepared?'

Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

- **1.** The domain of the function $\frac{x+1}{2x+1}$ is _____. (pp. 64–65)
- **2.** A function for which f(x) = f(-x) for all x in the domain of f is called an _____ function. (pp. 80–82)

Concepts and Vocabulary

- 5. The sine, cosine, cosecant, and secant functions have period ____; the tangent and cotangent functions have period
- 6. The domain of the tangent function is _____
- **7.** The range of the sine function is _____.

- **3.** True or False: The function $f(x) = \sqrt{x}$ is even. (pp. 80–82)
- 4. True or False: The equation $x^2 + 2x = (x + 1)^2 1$ is an identity. (p. 984)

Period is π .

- 8. True or False: The only even trigonometric functions are the cosine and secant functions.
- 9. True or False: All the trigonometric functions are periodic, with period 2π .
- 10. True or False: The range of the secant function is the set of positive real numbers.

Skill Building

1

.

٩

In Problems 11–26, use the fact that the trigonometric functions are periodic to find the exact value of each expression. Do not use a calculator.

11. sin 405°	12. cos 420°	13. tan 405°	14. sin 390°	15. csc 450°	16. sec 540°
17. cot 390°	18. sec 420°	19. $\cos \frac{33\pi}{4}$	20. $\sin \frac{9\pi}{4}$	21. tan(21π)	22. $\csc \frac{9\pi}{2}$
23. $\sec \frac{17\pi}{4}$	24. $\cot \frac{17\pi}{4}$	25. $\tan \frac{19\pi}{6}$	26. $\sec \frac{25\pi}{6}$		

In Problems 27–34, name the quadrant in which the angle θ lies.

$27. \sin \theta > 0,$	$\cos\theta < 0$	$28. \sin \theta < 0,$	$\cos \theta > 0$	29. $\sin \theta < 0$,	$\tan \theta < 0$
30. $\cos \theta > 0$,	$\tan \theta > 0$	31. $\cos \theta > 0$,	$\tan \theta < 0$	32. $\cos \theta < 0$,	$\tan \theta > 0$
33. sec $\theta < 0$,	$\sin \theta > 0$	34. $\csc \theta > 0$,	$\cos\theta < 0$		

In Problems 35–42, $\sin \theta$ and $\cos \theta$ are given. Find the exact value of each of the four remaining trigonometric functions.

35.
$$\sin \theta = -\frac{3}{5}, \ \cos \theta = \frac{4}{5}$$

36. $\sin \theta = \frac{4}{5}, \ \cos \theta = -\frac{3}{5}$
37. $\sin \theta = \frac{2\sqrt{5}}{5}, \ \cos \theta = \frac{\sqrt{5}}{5}$
38. $\sin \theta = -\frac{\sqrt{5}}{5}, \ \cos \theta = -\frac{2\sqrt{5}}{5}$
39. $\sin \theta = \frac{1}{2}, \ \cos \theta = \frac{\sqrt{3}}{2}$
40. $\sin \theta = \frac{\sqrt{3}}{2}, \ \cos \theta = \frac{1}{2}$
41. $\sin \theta = -\frac{1}{3}, \ \cos \theta = \frac{2\sqrt{2}}{3}$
42. $\sin \theta = \frac{2\sqrt{2}}{3}, \ \cos \theta = -\frac{1}{3}$

In Problems 43–58, find the exact value of each of the remaining trigonometric functions of θ .

43.
$$\sin \theta = \frac{12}{13}$$
, θ in quadrant II
 44. $\cos \theta = \frac{3}{5}$, θ in quadrant IV
 45. $\cos \theta = -\frac{4}{5}$, θ in quadrant III

 46. $\sin \theta = -\frac{5}{13}$, θ in quadrant III
 47. $\sin \theta = \frac{5}{13}$, $90^{\circ} < \theta < 180^{\circ}$
 48. $\cos \theta = \frac{4}{5}$, $270^{\circ} < \theta < 360^{\circ}$

 49. $\cos \theta = -\frac{1}{3}$, $\frac{\pi}{2} < \theta < \pi$
 50. $\sin \theta = -\frac{2}{3}$, $\pi < \theta < \frac{3\pi}{2}$
 51. $\sin \theta = \frac{2}{3}$, $\tan \theta < 0$

 52. $\cos \theta = -\frac{1}{4}$, $\tan \theta > 0$
 53. $\sec \theta = 2$, $\sin \theta < 0$
 54. $\csc \theta = 3$, $\cot \theta < 0$

 55. $\tan \theta = \frac{3}{4}$, $\sin \theta < 0$
 56. $\cot \theta = \frac{4}{3}$, $\cos \theta < 0$
 57. $\tan \theta = -\frac{1}{3}$, $\sin \theta > 0$

 58. $\sec \theta = -2$, $\tan \theta > 0$
 56. $\cot \theta = \frac{4}{3}$, $\cos \theta < 0$
 57. $\tan \theta = -\frac{1}{3}$, $\sin \theta > 0$

In Problems 59–76, use the even-odd properties to find the exact value of each expression. Do not use a calculator.

59. $\sin(-60^{\circ})$	60. $\cos(-30^{\circ})$	61. tan(-30°)	62. $\sin(-135^{\circ})$	63. $\sec(-60^{\circ})$	64. $\csc(-30^{\circ})$
65. sin(-90°)	66. $\cos(-270^{\circ})$	67. $\tan\left(-\frac{\pi}{4}\right)$	68. $\sin(-\pi)$	69. $\cos\left(-\frac{\pi}{4}\right)$	70. $\sin\left(-\frac{\pi}{3}\right)$
71. $tan(-\pi)$	72. $\sin\left(-\frac{3\pi}{2}\right)$	73. $\csc\left(-\frac{\pi}{4}\right)$	74. $sec(-\pi)$	75. $\operatorname{sec}\left(-\frac{\pi}{6}\right)$	76. $\csc\left(-\frac{\pi}{3}\right)$

In Problems 77–88, use properties of the trigonometric functions to find the exact value of each expression. Do not use a calculator.

77.
$$\sin^2 40^\circ + \cos^2 40^\circ$$
78. $\sec^2 18^\circ - \tan^2 18^\circ$ **79.** $\sin 80^\circ \csc 80^\circ$ **80.** $\tan 10^\circ \cot 10^\circ$ **81.** $\tan 40^\circ - \frac{\sin 40^\circ}{\cos 40^\circ}$ **82.** $\cot 20^\circ - \frac{\cos 20^\circ}{\sin 20^\circ}$ **83.** $\cos 400^\circ \cdot \sec 40^\circ$ **84.** $\tan 200^\circ \cdot \cot 20^\circ$ **85.** $\sin(-\frac{\pi}{12})\csc\frac{25\pi}{12}$ **86.** $\sec(-\frac{\pi}{18})\cdot\cos\frac{37\pi}{18}$ **87.** $\frac{\sin(-20^\circ)}{\cos 380^\circ} + \tan 200^\circ$ **88.** $\frac{\sin 70^\circ}{\cos(-430^\circ)} + \tan(-70^\circ)$

SECTION 5.3 Properties of the Trigonometric Functions 401

89. If $\sin \theta = 0.3$, find the value of:

$$\sin\theta + \sin(\theta + 2\pi) + \sin(\theta + 4\pi)$$

90. If $\cos \theta = 0.2$, find the value of:

$$\cos \theta + \cos(\theta + 2\pi) + \cos(\theta + 4\pi)$$

91. If $\tan \theta = 3$, find the value of:

$$\tan \theta + \tan(\theta + \pi) + \tan(\theta + 2\pi)$$

92. If $\cot \theta = -2$, find the value of:

$$\cot \theta + \cot(\theta - \pi) + \cot(\theta - 2\pi)$$

- 93. Find the exact value of: $\sin 1^\circ + \sin 2^\circ + \sin 3^\circ + \dots + \sin 358^\circ + \sin 359^\circ$
- 94. Find the exact value of: $\cos 1^\circ + \cos 2^\circ + \cos 3^\circ + \cdots + \cos 358^\circ + \cos 359^\circ$
- 95. What is the domain of the sine function?
- 96. What is the domain of the cosine function?
- 97. For what numbers θ is $f(\theta) = \tan \theta$ not defined?
 - **98.** For what numbers θ is $f(\theta) = \cot \theta$ not defined?
 - **99.** For what numbers θ is $f(\theta) = \sec \theta$ not defined?
- **100.** For what numbers θ is $f(\theta) = \csc \theta$ not defined?
- 101. What is the range of the sine function?
- **102.** What is the range of the cosine function?
- **103.** What is the range of the tangent function?
- **104.** What is the range of the cotangent function?
- **105.** What is the range of the secant function?
- **106.** What is the range of the cosecant function?
- **107.** Is the sine function even, odd, or neither? Is its graph symmetric? With respect to what?
- **108.** Is the cosine function even, odd, or neither? Is its graph symmetric? With respect to what?
- **109.** Is the tangent function even, odd, or neither? Is its graph symmetric? With respect to what?
- **110.** Is the cotangent function even, odd, or neither? Is its graph symmetric? With respect to what?
- **111.** Is the secant function even, odd, or neither? Is its graph symmetric? With respect to what?
- **112.** Is the cosecant function even, odd, or neither? Is its graph symmetric? With respect to what?

In Problems 113–118, use the periodic and even-odd properties. **113.** If $f(\theta) = \sin \theta$ and $f(a) = \frac{1}{3}$, find the exact value of: (a) f(-a) (b) $f(a) + f(a + 2\pi) + f(a + 4\pi)$ **114.** If $f(\theta) = \cos \theta$ and $f(a) = \frac{1}{4}$, find the exact value of: (a) f(-a) (b) $f(a) + f(a + 2\pi) + f(a - 2\pi)$


- 115. If f(θ) = tan θ and f(a) = 2, find the exact value of:
 (a) f(-a)
 (b) f(a) + f(a + π) + f(a + 2π)
- 116. If f(θ) = cot θ and f(a) = -3, find the exact value of:
 (a) f(-a)
 (b) f(a) + f(a + π) + f(a + 4π)
- 117. If f(θ) = sec θ and f(a) = -4, find the exact value of:
 (a) f(-a)
 (b) f(a) + f(a + 2π) + f(a + 4π)
- 118. If f(θ) = csc θ and f(a) = 2, find the exact value of:
 (a) f(-a)
 (b) f(a) + f(a + 2π) + f(a + 4π)
- **119.** Calculating the Time of a Trip From a parking lot, you want to walk to a house on the ocean. The house is located 1500 feet down a paved path that parallels the ocean, which is 500 feet away. See the illustration. Along the path you can walk 300 feet per minute, but in the sand on the beach you can only walk 100 feet per minute.

The time *T* to get from the parking lot to the beachhouse can be expressed as a function of the angle θ shown in the illustration and is

$$T(\theta) = 5 - \frac{5}{3\tan\theta} + \frac{5}{\sin\theta}, \qquad 0 < \theta < \frac{\pi}{2}$$

Calculate the time T if you walk directly from the parking lot to the house.

[**Hint:**
$$\tan \theta = \frac{500}{1500}$$
.]




120. Calculating the Time of a Trip Two oceanfront homes are located 8 miles apart on a straight stretch of beach, each a distance of 1 mile from a paved road that parallels the ocean. Sally can jog 8 miles per hour along the paved road, but only 3 miles per hour in the sand on the beach. Because of a river directly between the two houses, it is necessary to jog in the sand to the road, continue on the road, and then jog directly back in the sand to get from one house to the

other. See the illustration. The time T to get from one house to the other as a function of the angle θ shown in the illustration is

$$T(\theta) = 1 + \frac{2}{3\sin\theta} - \frac{1}{4\tan\theta} \qquad 0 < \theta < \frac{\pi}{2}$$

- (a) Calculate the time T for $\tan \theta = \frac{1}{4}$.
- (b) Describe the path taken.
- (c) Explain why θ must be larger than 14°.



- **Discussion and Writing**
- **132.** Write down five properties of the tangent function. Explain the meaning of each.
- **133.** Describe your understanding of the meaning of a periodic function.

'Are You Prepared?' Answers

1. $\left\{x \mid x \neq -\frac{1}{2}\right\}$ **2.** Even **3.** False **4.** True

- **121.** Show that the range of the tangent function is the set of all real numbers.
- **122.** Show that the range of the cotangent function is the set of all real numbers.
- **123.** Show that the period of $f(\theta) = \sin \theta$ is 2π .
 - **[Hint:** Assume that $0 exists so that <math>\sin(\theta + p) = \sin \theta$ for all θ . Let $\theta = 0$ to find p. Then

let
$$\theta = \frac{\pi}{2}$$
 to obtain a contradiction.]

- **124.** Show that the period of $f(\theta) = \cos \theta$ is 2π .
- **125.** Show that the period of $f(\theta) = \sec \theta$ is 2π .
- **126.** Show that the period of $f(\theta) = \csc \theta$ is 2π .
- **127.** Show that the period of $f(\theta) = \tan \theta$ is π .
- **128.** Show that the period of $f(\theta) = \cot \theta$ is π .
- **129.** Prove the reciprocal identities given in formula (2).
- **130.** Prove the quotient identities given in formula (3).
- **131.** Establish the identity:

$$(\sin\theta\cos\phi)^2 + (\sin\theta\sin\phi)^2 + \cos^2\theta = 1$$

- **134.** Explain how to find the value of sin 390° using periodic properties.
- **135.** Explain how to find the value of $\cos(-45^{\circ})$ using even-odd properties.

5.4 Graphs of the Sine and Cosine Functions*
 PREPARING FOR THIS SECTION Before getting started, review the following:
 Graphing Techniques: Transformations (Section 2.6, pp. 118–126)
 Now work the 'Are You Prepared?' problems on page 414.
 OBJECTIVES 1 Graph Transformations of the Sine Function

 Graph Transformations of the Cosine Function
 Graph Transformations of the Cosine Function
 Determine the Amplitude and Period of Sinusoidal Functions
 Graph Sinusoidal Functions Using Key Points
 Find an Equation for a Sinusoidal Graph

^{*}For those who wish to include phase shifts here, Section 5.6 can be covered immediately after Section 5.4 without loss of continuity.

Since we want to graph the trigonometric functions in the xy-plane, we shall use the traditional symbols x for the independent variable (or argument) and y for the dependent variable (or value at x) for each function. So we write the six trigonometric functions as

 $y = f(x) = \sin x$ $y = f(x) = \cos x$ $y = f(x) = \tan x$ $y = f(x) = \csc x$ $y = f(x) = \sec x$ $y = f(x) = \cot x$




Here the independent variable x represents an angle, measured in radians. In calculus, x will usually be treated as a real number. As we said earlier, these are equivalent ways of viewing x.

Graph Transformations of the Sine Function


Since the sine function has period 2π , we need to graph $y = \sin x$ only on the interval $[0, 2\pi]$. The remainder of the graph will consist of repetitions of this portion of the graph.

We begin by constructing Table 6, which lists some points on the graph of $y = \sin x, 0 \le x \le 2\pi$. As the table shows, the graph of $y = \sin x, 0 \le x \le 2\pi$, begins at the origin. As x increases from 0 to $\frac{\pi}{2}$, the value of $y = \sin x$ increases from 0 to 1; as x increases from $\frac{\pi}{2}$ to π to $\frac{3\pi}{2}$, the value of y decreases from 1 to 0 to -1; as x increases from $\frac{3\pi}{2}$ to 2π , the value of y increases from -1 to 0. If we plot the points listed in Table 6 and connect them with a smooth curve, we obtain the graph shown in Figure 49.

Figure 49



The graph in Figure 49 is one period, or cycle, of the graph of $y = \sin x$. To obtain a more complete graph of $y = \sin x$, we repeat this period in each direction, as shown in Figure 50(a). Figure 50(b) shows the graph on a TI-84 Plus graphing calculator.



(a)

Table 6

Figure 50

x	$y = \sin x$	(x, y)
0	0	(0, 0)
$\frac{\pi}{6}$	$\frac{1}{2}$	$\left(\frac{\pi}{6},\frac{1}{2}\right)$
$\frac{\pi}{2}$	1	$\left(\frac{\pi}{2},1\right)$
$\frac{5\pi}{6}$	$\frac{1}{2}$	$\left(\frac{5\pi}{6},\frac{1}{2}\right)$
π	0	(<i>π</i> , 0)
$\frac{7\pi}{6}$	$-\frac{1}{2}$	$\left(\frac{7\pi}{6},-\frac{1}{2}\right)$
$\frac{3\pi}{2}$	-1	$\left(\frac{3\pi}{2}, -1\right)$
$\frac{11\pi}{6}$	$-\frac{1}{2}$	$\left(\frac{11\pi}{6},-\frac{1}{2}\right)$
2π	0	(2 <i>π</i> , 0)

(b)

The graph of $y = \sin x$ illustrates some of the facts that we already know about the sine function.

Properties of the Sine Function

- 1. The domain is the set of all real numbers.
- **2.** The range consists of all real numbers from -1 to 1, inclusive.
- **3.** The sine function is an odd function, as the symmetry of the graph with respect to the origin indicates.
- **4.** The sine function is periodic, with period 2π .
- 5. The x-intercepts are \ldots , -2π , $-\pi$, 0, π , 2π , 3π , \ldots ; the y-intercept is 0.
- 6. The maximum value is 1 and occurs at $x = \dots, -\frac{3\pi}{2}, \frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \dots;$

the minimum value is -1 and occurs at $x = \dots, -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}, \frac{11\pi}{2}, \dots$

NOW WORK PROBLEMS 9, 11, AND 13.

The graphing techniques introduced in Chapter 2, Section 2.6, may be used to graph functions that are transformations of the sine function.


EXAMPLE 1 Graphing Variations of y = sin x Using Transformations

Use the graph of
$$y = \sin x$$
 to graph $y = \sin\left(x - \frac{\pi}{4}\right)$.

Solution


Figure 51 illustrates the steps.






CHECK: Figure 52 shows the graph of $Y_1 = \sin\left(x - \frac{\pi}{4}\right)$.

Figure 52



EXAMPLE 2 Graphing Variations of y = sin x Using Transformations

Use the graph of $y = \sin x$ to graph $y = -\sin x + 2$.



✓ CHECK: Figure 54 shows the graph of $Y_1 = -\sin x + 2$.





Table 7


x	$y = \cos x$	(x, y)
0	1	(0, 1)
$\frac{\pi}{3}$	$\frac{1}{2}$	$\left(\frac{\pi}{3},\frac{1}{2}\right)$
$\frac{\pi}{2}$	0	$\left(\frac{\pi}{2}, 0\right)$
$\frac{2\pi}{3}$	$-\frac{1}{2}$	$\left(\frac{2\pi}{3},-\frac{1}{2}\right)$
π	-1	(<i>π</i> , −1)
$\frac{4\pi}{3}$	$-\frac{1}{2}$	$\left(\frac{4\pi}{3},-\frac{1}{2}\right)$
$\frac{3\pi}{2}$	0	$\left(\frac{3\pi}{2},0\right)$
$\frac{5\pi}{3}$	$\frac{1}{2}$	$\left(\frac{5\pi}{3},\frac{1}{2}\right)$
2π	1	(2 <i>π</i> , 1)

Graph Transformations of the Cosine Function

The cosine function also has period 2π . We proceed as we did with the sine function by constructing Table 7, which lists some points on the graph of $y = \cos x$, $0 \le x \le 2\pi$. As the table shows, the graph of $y = \cos x$, $0 \le x \le 2\pi$, begins at the point (0, 1). As x increases from 0 to $\frac{\pi}{2}$ to π , the value of y decreases from 1 to 0 to -1; as x increases from π to $\frac{3\pi}{2}$ to 2π , the value of y increases from -1 to 0 to 1. As before, we plot the points in Table 7 to get one period or cycle of the graph. See Figure 55.



A more complete graph of $y = \cos x$ is obtained by repeating this period in each direction, as shown in Figure 56(a). Figure 56(b) shows the graph on a TI-84 Plus graphing calculator.



The graph of $y = \cos x$ illustrates some of the facts that we already know about the cosine function.

Properties of the Cosine Function

- 1. The domain is the set of all real numbers.
- **2.** The range consists of all real numbers from -1 to 1, inclusive.
- **3.** The cosine function is an even function, as the symmetry of the graph with respect to the *y*-axis indicates.
- 4. The cosine function is periodic, with period 2π .
- 5. The x-intercepts are $\ldots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \ldots$; the y-intercept is 1.
- 6. The maximum value is 1 and occurs at $x = \dots, -2\pi, 0, 2\pi, 4\pi, 6\pi, \dots$; the minimum value is -1 and occurs at $x = \dots, -\pi, \pi, 3\pi, 5\pi, \dots$


Again, the graphing techniques from Chapter 2 may be used to graph transformations of the cosine function.

EXAMPLE 3 Graphing Variations of y = cos x Using Transformations

Use the graph of $y = \cos x$ to graph $y = 2 \cos x$.


Solution Figure 57 illustrates the steps.

Figure 57



3π

✓ **CHECK:** Figure 58 shows the graph $Y_1 = 2 \cos x$.




EXAMPLE 4 Graphing Variations of y = cos x Using Transformations


Use the graph of $y = \cos x$ to graph $y = \cos(3x)$.

Solution Figure 59 illustrates the graph, which is a horizontal compression of the graph of $y = \cos x$. (Multiply each x-coordinate by $\frac{1}{3}$.) Notice that, due to this compression, the period of $y = \cos(3x)$ is $\frac{2\pi}{3}$, whereas the period of $y = \cos x$ is 2π .





✓ **CHECK:** Figure 60 shows the graph of $Y_1 = \cos(3x)$.




NOW WORK PROBLEM 33.


Sinusoidal Graphs

Shift the graph of $y = \cos x$ to the right $\frac{\pi}{2}$ units to obtain the graph of $y = \cos\left(x - \frac{\pi}{2}\right)$. See Figure 61(a). Now look at the graph of $y = \sin x$ in Figure 61(b). We see that the graph of $y = \sin x$ is the same as the graph of $y = \cos\left(x - \frac{\pi}{2}\right)$.





Based on Figure 61, we conjecture that




(We shall prove this fact in Chapter 6.) Because of this similarity, the graphs of sine functions and cosine functions are referred to as **sinusoidal graphs**. Let's look at some general properties of sinusoidal graphs.

 $\sin x = \cos\left(x - \frac{\pi}{2}\right)$

Determine the Amplitude and Period of Sinusoidal Functions

In Example 3 we obtained the graph of $y = 2 \cos x$, which we reproduce in Figure 62. Notice that the values of $y = 2 \cos x$ lie between -2 and 2, inclusive.




In general, the values of the functions $y = A \sin x$ and $y = A \cos x$, where $A \neq 0$, will always satisfy the inequalities


$$-|A| \le A \sin x \le |A|$$
 and $-|A| \le A \cos x \le |A|$

respectively. The number |A| is called the **amplitude** of $y = A \sin x$ or $y = A \cos x$. See Figure 63.

Figure 63




In Example 4, we obtained the graph of $y = \cos(3x)$, which we reproduce in Figure 64. Notice that the period of this function is $\frac{2\pi}{3}$.



In general, if $\omega > 0$, the functions $y = \sin(\omega x)$ and $y = \cos(\omega x)$ will have period $T = \frac{2\pi}{\omega}$. To see why, recall that the graph of $y = \sin(\omega x)$ is obtained from the graph of $y = \sin x$ by performing a horizontal compression or stretch by a factor $\frac{1}{\omega}$. This horizontal compression replaces the interval $[0, 2\pi]$, which contains one period of the graph of $y = \sin x$, by the interval $\left[0, \frac{2\pi}{\omega}\right]$, which contains one period of the graph of $y = \sin(\omega x)$. The period of the functions $y = \sin(\omega x)$ and $y = \cos(\omega x), \omega > 0$, is $\frac{2\pi}{\omega}$.

For example, for the function $y = \cos(3x)$, graphed in Figure 64, $\omega = 3$, so the period is $\frac{2\pi}{\omega} = \frac{2\pi}{3}$.

One period of the graph of $y = \sin(\omega x)$ or $y = \cos(\omega x)$ is called a **cycle**. Figure 65 illustrates the general situation. The blue portion of the graph is one cycle.



If $\omega < 0$ in $y = \sin(\omega x)$ or $y = \cos(\omega x)$, we use the Even–Odd Properties of the sine and cosine functions as follows:

 $\sin(-\omega x) = -\sin(\omega x)$ and $\cos(-\omega x) = \cos(\omega x)$

This gives us an equivalent form in which the coefficient of x is positive. For example,

$$\sin(-2x) = -\sin(2x)$$
 and $\cos(-\pi x) = \cos(\pi x)$

Theorem

If $\omega > 0$, the amplitude and period of $y = A \sin(\omega x)$ and $y = A \cos(\omega x)$ are given by

Amplitude =
$$|A|$$
 Period = $T = \frac{2\pi}{\omega}$ (1)

EXAMPLE 5 Finding the Amplitude and Period of a Sinusoidal Function

Determine the amplitude and period of $y = 3 \sin(4x)$.

Solution Comparing $y = 3\sin(4x)$ to $y = A\sin(\omega x)$, we find that A = 3 and $\omega = 4$. From equation (1),

Amplitude =
$$|A| = 3$$
 Period = $T = \frac{2\pi}{\omega} = \frac{2\pi}{4} = \frac{\pi}{2}$

4 Graph Sinusoidal Functions Using Key Points

Earlier, we graphed sine and cosine functions using tranformations. We now introduce another method that can be used to graph these functions.


Figure 66 shows one cycle of the graphs of $y = \sin x$ and $y = \cos x$ on the interval $[0, 2\pi]$. Notice that each graph consists of four parts corresponding to the four subintervals:

$$\left[0,\frac{\pi}{2}\right], \left[\frac{\pi}{2},\pi\right], \left[\pi,\frac{3\pi}{2}\right], \left[\frac{3\pi}{2},2\pi\right]$$

Each subinterval is of length $\frac{\pi}{2}$ (the period 2π divided by 4, the number of parts), and the endpoints of these intervals give rise to five key points on the graph:

For
$$y = \sin x$$
: $(0, 0)$, $\left(\frac{\pi}{2}, 1\right)$, $(\pi, 0)$, $\left(\frac{3\pi}{2}, -1\right)$, $(2\pi, 0)$
For $y = \cos x$: $(0, 1)$, $\left(\frac{\pi}{2}, 0\right)$, $(\pi, -1)$, $\left(\frac{3\pi}{2}, 0\right)$, $(2\pi, 1)$

See Figure 66.



When graphing a sinusoidal function of the form $y = A \sin(\omega x)$ or $y = A \cos(\omega x)$ we use the amplitude to determine the maximum and minimum values of the function. The period is used to divide the *x*-axis into four subintervals. The endpoints of the subintervals give rise to five key points on the graph, which are used to sketch one cycle. Finally, extend the graph in either direction to make it complete.

Let's look at an example.

EXAMPLE 6 Graphing a Sinusoidal Function Using Key Points

Graph: $y = 3\sin(4x)$

Solution

From Example 5, the amplitude is 3 and the period is $\frac{\pi}{2}$. Because the amplitude is 3, the graph of $y = 3 \sin(4x)$ will lie between -3 and 3 on the y-axis. Because the period is $\frac{\pi}{2}$, one cycle will begin at x = 0 and end at $x = \frac{\pi}{2}$. We divide the interval $\left[0, \frac{\pi}{2}\right]$ into four subintervals, each of length $\frac{\pi}{2} \div 4 = \frac{\pi}{8}$: $\left[0, \frac{\pi}{8}\right], \left[\frac{\pi}{8}, \frac{\pi}{4}\right], \left[\frac{\pi}{4}, \frac{3\pi}{8}\right], \left[\frac{3\pi}{8}, \frac{\pi}{2}\right]$

NOTE


We could also obtain the five key points by evaluating $y = 3 \sin(4x)$ at the endopoints of each subinterval.

The endpoints of these intervals give rise to the *x*-coordinates of the five key points on the graph. To obtain the *y*-coordinates of the five key points, multiply the *y*-coordinates of the five key points in $y = \sin x$ by A = 3. Refer to Figure 66. The five key points are

$$(0,0), \quad \left(\frac{\pi}{8},3\right), \quad \left(\frac{\pi}{4},0\right), \quad \left(\frac{3\pi}{8},-3\right), \quad \left(\frac{\pi}{2},0\right)$$

We plot these five points and fill in the graph of the sine curve as shown in Figure 67(a). We extend the graph in either direction to obtain the complete graph shown in Figure 67(b).

To graph a sinusoidal function using a graphing utility, we use the amplitude to set Ymin and Ymax and use the period to set Xmin and Xmax. Figure 67(c) shows the graph using a graphing utility.



CHECK: Graph $y = 3 \sin(4x)$ by hand using transformations. Which graphing method do you prefer?

NOW WORK PROBLEM 47.

EXAMPLE 7 Finding the Amplitude and Period of a Sinusoidal Function and Graphing It Using Key Points

Determine the amplitude and period of $y = -4\cos(\pi x)$, and graph the function.

Solution

Comparing $y = -4\cos(\pi x)$ with $y = A\cos(\omega x)$, we find that A = -4 and $\omega = \pi$. The amplitude is |A| = |-4| = 4, and the period is $T = \frac{2\pi}{\omega} = \frac{2\pi}{\pi} = 2$.

The graph of $y = -4 \cos(\pi x)$ will lie between -4 and 4 on the y-axis. One cycle will begin at x = 0 and end at x = 2. We divide the interval [0, 2] into four subintervals, each of length $2 \div 4 = \frac{1}{2}$:


$$\left[0,\frac{1}{2}\right], \quad \left[\frac{1}{2},1\right], \quad \left[1,\frac{3}{2}\right], \quad \left[\frac{3}{2},2\right].$$

The five key points on the graph are

$$(0, -4), (\frac{1}{2}, 0), (1, 4), (\frac{3}{2}, 0), (2, -4).$$

We plot these five points and fill in the graph of the cosine function as shown in Figure 68(a). Extending the graph in either direction, we obtain Figure 68(b).

Figure 68(c) shows the graph using a graphing utility.



✓ **CHECK:** Graph $y = -4 \cos(\pi x)$ by hand using transformations. Which graphing method do you prefer?

EXAMPLE 8 Finding the Amplitude and Period of a Sinusoidal Function and Graphing It Using Key Points

Determine the amplitude and period of $y = 2\sin\left(-\frac{\pi}{2}x\right)$, and graph the function.

Solution

$$y = -2\sin\!\left(\frac{\pi}{2}x\right)$$

Comparing
$$y = -2\sin\left(\frac{\pi}{2}x\right)$$
 to $y = A\sin(\omega x)$, we find that $A = -2$ and $\omega = \frac{\pi}{2}$.

The amplitude is |A| = 2, and the period is $T = \frac{2\pi}{\omega} = \frac{2\pi}{\frac{\pi}{2}} = 4$.


Since the sine function is odd, we can use the equivalent form:

The graph of $y = -2 \sin\left(\frac{\pi}{2}x\right)$ will lie between -2 and 2 on the y-axis. One cycle will begin at x = 0 and end at x = 4. We divide the interval [0, 4] into four subintervals, each of length $4 \div 4 = 1$:

The five key points on the graph are

(0,0), (1,-2), (2,0), (3,2), (4,0)

We plot these five points and fill in the graph of the sine function as shown in Figure 69(a). Extending the graph in either direction, we obtain Figure 69(b). Figure 69(c) shows the graph using a graphing utility.




We can also use the ideas of amplitude and period to identify a sinusoidal function when its graph is given.

EXAMPLE 9 Finding an Equation for a Sinusoidal Graph

Find an equation for the graph shown in Figure 70.

Figure 70



Solution

The graph has the characteristics of a cosine function. Do you see why? So we view the equation as a cosine function $y = A \cos(\omega x)$ with A = 3 and period T = 1. Then $\frac{2\pi}{\omega} = 1$, so $\omega = 2\pi$. The cosine function whose graph is given in Figure 70 is $y = A \cos(\omega x) = 3 \cos(2\pi x)$

✓ CHECK: Graph $Y_1 = 3\cos(2\pi x)$ and compare the result with Figure 70.

EXAMPLE 10 Finding an Equation for a Sinusoidal Graph

Find an equation for the graph shown in Figure 71.

Solution

-2 -3 5

Figure 71

The graph is sinusoidal, with amplitude |A| = 2. The period is 4, so $\frac{2\pi}{\omega} = 4$ or $\omega = \frac{\pi}{2}$. Since the graph passes through the origin, it is easiest to view the equation as a sine function,^{*} but notice that the graph is actually the reflection of a sine function about the *x*-axis (since the graph is decreasing near the origin). Thus, we have

$$y = -A\sin(\omega x) = -2\sin\left(\frac{\pi}{2}x\right)$$

CHECK: Graph $Y_1 = -2\sin\left(\frac{\pi}{2}x\right)$ and compare the result with Figure 71.

NOW WORK PROBLEMS 71 AND 75.

^{*}The equation could also be viewed as a cosine function with a horizontal shift, but viewing it as a sine function is easier.

5.4 Assess Your Understanding

'Are You Prepared?'

Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

- **1.** Use transformations to graph $y = 3x^2$ (pp. 120–122)
- **2.** Use transformations to graph $y = -x^2$. (pp. 123–124)

Concepts and Vocabulary

- 3. The maximum value of $y = \sin x$ is _____ and occurs at x =____.
- **4.** The function $y = A \sin(\omega x)$, A > 0, has amplitude 3 and period 2; then $A = _$ and $\omega = _$.
- **5.** The function $y = 3\cos(6x)$ has amplitude _____ and period
- **6.** *True or False:* The graphs of $y = \sin x$ and $y = \cos x$ are identical except for a horizontal shift.
- 7. *True or False:* For $y = 2\sin(\pi x)$, the amplitude is 2 and the period is $\frac{\pi}{2}$.
- **8.** *True or False:* The graph of the sine function has infinitely many *x*-intercepts.

Skill Building

In Problems 9–18, if necessary, refer to the graphs to answer each question.


- 9. What is the *y*-intercept of $y = \sin x$?
- 11. For what numbers $x, -\pi \le x \le \pi$, is the graph of $y = \sin x$ increasing?
- **10.** What is the *y*-intercept of $y = \cos x$?
- **12.** For what numbers x, $-\pi \le x \le \pi$, is the graph of $y = \cos x$ decreasing?


- 13. What is the largest value of $y = \sin x$?
 - **15.** For what numbers $x, 0 \le x \le 2\pi$, does sin x = 0?
 - **17.** For what numbers $x, -2\pi \le x \le 2\pi$, does sin x = 1? What about sin x = -1?
- **14.** What is the smallest value of $y = \cos x$?

C. $y = \sin\left(x - \frac{\pi}{2}\right)$

- **16.** For what numbers $x, 0 \le x \le 2\pi$, does $\cos x = 0$?
- **18.** For what numbers $x, -2\pi \le x \le 2\pi$, does $\cos x = 1$? What about $\cos x = -1$?

In Problems 19 and 20, match the graph to a function. Three answers are possible.





In Problems 21–36, use transformations to graph each function. Verify your results using a graphing utility.


21. $y = 3 \sin x$	22. $y = 4 \cos x$	23. $y = -\cos x$	24. $y = -\sin x$
25. $y = \sin x - 1$	26. $y = \cos x + 1$	27. $y = \sin(x - \pi)$	28. $y = \cos(x + \pi)$
29. $y = \sin(\pi x)$	30. $y = \cos\left(\frac{\pi}{2}x\right)$	31. $y = 2 \sin x + 2$	32. $y = 3\cos x + 3$
33. $y = 4\cos(2x)$	34. $y = 3\sin\left(\frac{1}{2}x\right)$	35. $y = -2\sin x + 2$	36. $y = -3\cos x - 2$


In Problems 37–46, determine the amplitude and period of each function without graphing.

37.
$$y = 2 \sin x$$


38. $y = 3 \cos x$
39. $y = -4 \cos(2x)$
40. $y = -\sin(\frac{1}{2}x)$
41. $y = 6 \sin(\pi x)$
42. $y = -3 \cos(3x)$
43. $y = -\frac{1}{2}\cos(\frac{3}{2}x)$
44. $y = \frac{4}{3}\sin(\frac{2}{3}x)$
45. $y = \frac{5}{3}\sin(-\frac{2\pi}{3}x)$
46. $y = \frac{9}{5}\cos(-\frac{3\pi}{2}x)$


In Problems 47–56, match the given function to one of the graphs (A)–(J).








In Problems 57–60, match the given function to one of the graphs (A)-(D).




















Applications and Extensions

89. Alternating Current (ac) Circuits The current *I*, in amperes, flowing through an ac (alternating current) circuit at time *t* is


$$I = 220\sin(60\pi t), \qquad t \ge 0$$

What is the period? What is the amplitude? Graph this function over two periods.

90. Alternating Current (ac) Circuits The current *I*, in amperes, flowing through an ac (alternating current) circuit at time *t* is

$$I = 120\sin(30\pi t), \qquad t \ge 0$$

What is the period? What is the amplitude? Graph this function over two periods.



91. Alternating Current (ac) Generators The voltage V produced by an ac generator is

 $V = 220\sin(120\pi t)$

π

- (a) What is the amplitude? What is the period?
- (b) Graph V over two periods, beginning at t = 0.
- (c) If a resistance of R = 10 ohms is present, what is the current *I*?
- [**Hint:** Use Ohm's Law, V = IR.]
- (d) What is the amplitude and period of the current *I*?
- (e) Graph I over two periods, beginning at t = 0.
- **92. Alternating Current (ac) Generators** The voltage V produced by an ac generator is

$$V = 120\sin(120\pi t)$$


418 CHAPTER 5 Trigonometric Functions

- (a) What is the amplitude? What is the period?
- (b) Graph V over two periods, beginning at t = 0.
- (c) If a resistance of R = 20 ohms is present, what is the current *I*?
 - [**Hint:** Use Ohm's Law, V = IR.]
- (d) What is the amplitude and period of the current *I*?
- (e) Graph *I* over two periods, beginning at t = 0.
- **93.** Alternating Current (ac) Generators The voltage V produced by an ac generator is sinusoidal. As a function of time, the voltage V is

$$V = V_0 \sin(2\pi f t)$$

where f is the **frequency**, the number of complete oscillations (cycles) per second. [In the United States and Canada, f is 60 hertz (Hz).] The **power** P delivered to a resistance Rat any time t is defined as

$$P = \frac{V^2}{R}$$



Power in an ac generator

(a) Show that
$$P = \frac{V_0^2}{R} \sin^2(2\pi ft)$$
.

- (b) The graph of P is shown in the figure. Express P as a sinusoidal function.
- (c) Deduce that


$$\sin^2(2\pi ft) = \frac{1}{2} [1 - \cos(4\pi ft)]$$


Discussion and Writing

- **98.** Explain how you would scale the *x*-axis and *y*-axis before graphing $y = 3\cos(\pi x)$.
- **99.** Explain the term *amplitude* as it relates to the graph of a sinusoidal function.

'Are You Prepared?' Answers

1. Vertical stretch by a factor of 3






94. Biorhythms In the theory of biorhythms, a sine function of the form

$$P = 50\sin(\omega t) + 50$$

is used to measure the percent P of a person's potential at time t, where t is measured in days and t = 0 is the person's birthday. Three characteristics are commonly measured:

Physical potential: period of 23 days Emotional potential: period of 28 days Intellectual potential: period of 33 days

- (a) Find ω for each characteristic.
- (b) Graph all three functions.
- (c) Is there a time t when all three characteristics have 100% potential? When is it?
- (d) Suppose that you are 20 years old today (t = 7305 days). Describe your physical, emotional, and intellectual potential for the next 30 days.



- **95.** Graph $y = |\cos x|, -2\pi \le x \le 2\pi$.
- **96.** Graph $y = |\sin x|, -2\pi \le x \le 2\pi$.
- **97.** Draw a quick sketch of $y = \sin x$. Be sure to label at least five points.
- **100.** Explain how the amplitude and period of a sinusoidal graph are used to establish the scale on each coordinate axis.
- **101.** Find an application in your major field that leads to a sinusoidal graph. Write a paper about your findings.

5.5 Graphs of the Tangent, Cotangent, Cosecant, and Secant Functions

PREPARING FOR THIS SECTION Before getting started, review the following:

• Vertical Asymptotes (Section 3.3 pp. 189–191)


Now work the 'Are You Prepared?' problems on page 423.

OBJECTIVES 1 Graph Transformations of the Tangent Function and Cotangent Function

2 Graph Transformations of the Cosecant Function and Secant Function

Graphs Transformations of the Tangent Function and Cotangent Function

Because the tangent function has period π , we only need to determine the graph over some interval of length π . The rest of the graph will consist of repetitions of that graph. Because the tangent function is not defined at $\dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$, we will concentrate on the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, of length π , and construct Table 8, which lists some points on the graph of $y = \tan x, -\frac{\pi}{2} < x < \frac{\pi}{2}$. We plot the points in the table and connect them with a smooth curve. See Figure 72 for a partial graph of $y = \tan x$, where $-\frac{\pi}{3} \le x \le \frac{\pi}{3}$.





	X	Y1			
<u>π</u> 2 - <u>π</u> 2	1.5 1.57 ↑ 1.5707 ↑ 1.5708 ↑ 1.57 ↑ 1.57 ↑ 1.571	14.101 1255.8 10381 ERROR -14.1 -1256 ERROR			
-	YiBtan(X)				

To complete one period of the graph of $y = \tan x$, we need to investigate the behavior of the function as x approaches $-\frac{\pi}{2}$ and $\frac{\pi}{2}$. We must be careful, though, because $y = \tan x$ is not defined at these numbers. To determine this behavior, we use the table feature on a graphing utility.

See Table 9. As x gets closer to $\frac{\pi}{2} \approx 1.5708$, but remains less than $\frac{\pi}{2}$, the values of tan x are positive and getting larger, so tan x approaches $\infty \left(\lim_{x \to \frac{\pi}{2}} \tan x = \infty\right)$.


In other words, the vertical line $x = \frac{\pi}{2}$ is a vertical asymptote to the graph of $y = \tan x$. As x gets closer to $-\frac{\pi}{2}$, but remains greater than $-\frac{\pi}{2}$, the values of $\tan x$ are negative and getting larger in magnitude, so $\tan x$ approaches $-\infty$ $\left(\lim_{x \to -\frac{\pi}{2}+\frac{\pi}{2}} \tan x = -\infty\right)$. In other words, the vertical line $x = -\frac{\pi}{2}$ is also a vertical asymptote to the graph

vertical asymptote to the graph.

With these observations, we can complete one period of the graph. We obtain the complete graph of $y = \tan x$ by repeating this period, as shown in Figure 73(a). Figure 73(b) shows the graph of $y = \tan x, -\infty < x < \infty$, using a graphing

utility. Notice we used dot mode when graphing $y = \tan x$, $-\infty < x < \infty$, using a graphing utility. Notice we used dot mode when graphing $y = \tan x$. Do you know why?






The graph of $y = \tan x$ illustrates some facts that we already know about the tangent function.

Properties of the Tangent Function


- **1.** The domain is the set of all real numbers, except odd multiples of $\frac{\pi}{2}$.
- 2. The range is the set of all real numbers.
- **3.** The tangent function is an odd function, as the symmetry of the graph with respect to the origin indicates.
- 4. The tangent function is periodic, with period π .
- 5. The x-intercepts are $\ldots, -2\pi, -\pi, 0, \pi, 2\pi, 3\pi, \ldots$; the y-intercept is 0.
- 6. Vertical asymptotes occur at $x = \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$

Graph: $y = 2 \tan x$

Solution We start with the graph of $y = \tan x$ and vertically stretch it by a factor of 2. See Figure 74.



CHECK: Figure 75 shows the graph using a graphing utility.



EXAMPLE 2


Graphing Variations of $y = \tan x$ Using Transformations

Graph:
$$y = -\tan\left(x + \frac{\pi}{4}\right)$$

Solution

We start with the graph of $y = \tan x$. See Figure 76.





✓ CHECK: Graph $Y_1 = -\tan\left(x + \frac{\pi}{4}\right)$ and compare the result to Figure 76(c). ◀

NOW WORK PROBLEM 25.



x	$y = \cot x$	(x, y)
$\frac{\pi}{6}$	$\sqrt{3}$	$\left(\frac{\pi}{6},\sqrt{3}\right)$
$\frac{\pi}{4}$	1	$\left(\frac{\pi}{4},1\right)$
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{3}$	$\left(\frac{\pi}{3},\frac{\sqrt{3}}{3}\right)$
$\frac{\pi}{2}$	0	$\left(\frac{\pi}{2},0\right)$
$\frac{2\pi}{3}$	$-\frac{\sqrt{3}}{3}$	$\left(\frac{2\pi}{3},-\frac{\sqrt{3}}{3}\right)$
$\frac{3\pi}{4}$	-1	$\left(\frac{3\pi}{4},-1\right)$
$\frac{5\pi}{6}$	$-\sqrt{3}$	$\left(\frac{5\pi}{6}, -\sqrt{3}\right)$



Χ	Y1			
5,11,000 1,000 1,000 1,000 1,14 1,44 1,44	1.8305 9.9666 99.997 1000 -7.015 -24.03 -627.9			
Yı∎1∕tan(X)				


We obtain the graph of $y = \cot x$ as we did the graph of $y = \tan x$. The period of $y = \cot x$ is π . Because the cotangent function is not defined for integer multiples of π , we will concentrate on the interval $(0, \pi)$. Table 10 lists some points on the graph of $y = \cot x$, $0 < x < \pi$.

See Table 11. As x approaches 0, but remains greater than 0, the values of cot x will be positive and large; so as x approaches 0, with x > 0, cot x approaches ∞

 $\left(\lim_{x\to 0^+} \cot x = \infty\right)$. Similarly, as x approaches π , but remains less than π , the values of $\cot x$ will be negative and will approach $-\infty\left(\lim_{x\to\pi^-} \cot x = -\infty\right)$. Figure 77 shows the graph.

Figure 77

 $y = \cot x, -\infty < x < \infty, x$ not equal to integer multiples of $\pi, -\infty < y < \infty$




Graph Transformations of the Cosecant Function and Secant Function

The cosecant and secant functions, sometimes referred to as **reciprocal functions**, are graphed by making use of the reciprocal identities

$$\csc x = \frac{1}{\sin x}$$
 and $\sec x = \frac{1}{\cos x}$

For example, the value of the cosecant function $y = \csc x$ at a given number x equals the reciprocal of the corresponding value of the sine function, provided that the value of the sine function is not 0. If the value of sin x is 0, then x is an integer multiple of π . At such numbers, the cosecant function is not defined. In fact, the graph of the cosecant function has vertical asymptotes at integer multiples of π . Figure 78 shows the graph.



EXAMPLE 3 Graphing Variations of y = csc x Using Transformations

Graph: $y = 2 \csc x - 1$






Figure 79



CHECK: Graph $Y_1 = 2 \csc x - 1$ and compare the result with Figure 79.

NOW WORK PROBLEM 37.

Using the idea of reciprocals, we can similarly obtain the graph of $y = \sec x$. See Figure 80.



5.5 Assess Your Understanding

'Are You Prepared?'

Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

- 1. The graph of $y = \frac{3x-6}{x-4}$ has a vertical asymptote. What is it? (pp. 189–191)
- **2.** *True or False:* If a function f has the vertical asymptote x = c, then f(c) is not defined. (pp. 189–191)

Concepts and Vocabulary

- 3. The graph of y = tan x is symmetric with respect to the _____ and has vertical asymptotes at _____.
- 4. The graph of y = sec x is symmetric with respect to the _____ and has vertical asymptotes at _____.


Skill Building


In Problems 7–16, if necessary, refer to the graphs to answer each question.

- 7. What is the y-intercept of $y = \tan x$?
- 9. What is the *y*-intercept of $y = \sec x$?
- **11.** For what numbers $x, -2\pi \le x \le 2\pi$, does sec x = 1? What about sec x = -1?
- **13.** For what numbers $x, -2\pi \le x \le 2\pi$, does the graph of $y = \sec x$ have vertical asymptotes?
- **15.** For what numbers $x, -2\pi \le x \le 2\pi$, does the graph of $y = \tan x$ have vertical asymptotes?

In Problems 17–20, match each function to its graph.

- 5. It is easiest to graph $y = \sec x$ by first sketching the graph of _____.
- 6. *True or False:* The graphs of $y = \tan x$, $y = \cot x$, $y = \sec x$, and $y = \csc x$ each have infinitely many vertical asymptotes.
- 8. What is the *y*-intercept of $y = \cot x$?
- **10.** What is the *y*-intercept of $y = \csc x$?
- **12.** For what numbers $x, -2\pi \le x \le 2\pi$, does $\csc x = 1$? What about $\csc x = -1$?
- 14. For what numbers $x, -2\pi \le x \le 2\pi$, does the graph of $y = \csc x$ have vertical asymptotes?
- **16.** For what numbers $x, -2\pi \le x \le 2\pi$, does the graph of $y = \cot x$ have vertical asymptotes?






In Problems 21-44, use tranformations to graph each function. Verify your results using a graphing utility.

21. $y = -\sec x$	22. $y = -\cot x$	$23. \ y = \sec\left(x - \frac{\pi}{2}\right)$	24. $y = \csc(x - \pi)$
25. $y = \tan(x - \pi)$	26. $y = \cot(x - \pi)$	27. $y = 3 \tan(2x)$	28. $y = 4 \tan\left(\frac{1}{2}x\right)$
29. $y = \sec(2x)$	30. $y = \csc\left(\frac{1}{2}x\right)$	31. $y = \cot(\pi x)$	32. $y = \cot(2x)$
33. $y = -3 \tan(4x)$	34. $y = -3 \tan(2x)$	$35. \ y = 2 \sec\left(\frac{1}{2}x\right)$	36. $y = 2 \sec(3x)$
37. $y = -3 \csc\left(x + \frac{\pi}{4}\right)$	$38. \ y = -2 \tan\left(x + \frac{\pi}{4}\right)$	$39. \ y = \frac{1}{2} \cot\left(x - \frac{\pi}{4}\right)$	$40. \ y = 3 \sec\left(x + \frac{\pi}{2}\right)$
41. $y = \tan x + 2$	42. $y = \cot x - 1$	43. $y = \sec\left(x + \frac{\pi}{2}\right) - 1$	44. $y = \csc\left(x - \frac{\pi}{4}\right) + 1$

Applications and Extensions


45. Carrying a Ladder around a Corner Two hallways, one of width 3 feet, the other of width 4 feet, meet at a right angle. See the illustration.



(a) Show that the length L of the line segment shown as a function of the angle θ is

$$L(\theta) = 3 \sec \theta + 4 \csc \theta$$

- (b) Graph $L = L(\theta), 0 < \theta < \frac{\pi}{2}$.
- (c) For what value of θ is *L* the least?
- (d) What is the length of the longest ladder that can be carried around the corner? Why is this also the least value of *L*?
- **46. A Rotating Beacon** Suppose that a fire truck is parked in front of a building as shown in the figure.



The beacon light on top of the fire truck is located 10 feet from the wall and has a light on each side. If the beacon light rotates 1 revolution every 2 seconds, then a model for determining the distance d that the beacon of light is from point A on the wall after t seconds is given by

$$d = 10 \tan(\pi t)$$

- (a) Graph $d = 10 \tan(\pi t)$ for $0 \le t \le 2$.
- (b) For what values of *t* is the function undefined? Explain what this means in terms of the beam of light on the wall.(c) Fill in the following table.

t	$d = 10 \tan(\pi t)$
0	
0.1	
0.2	
0.3	
0.4	

- (d) Compute $\frac{d(0.1) d(0)}{0.1 0}$, $\frac{d(0.2) d(0.1)}{0.2 0.1}$, and so on, for each consecutive value of *t*. These are called **first differences**.
- (e) Interpret the first differences found in part (d). What is happening to the speed of the beam of light as *d* increases?
- 47. Exploration Graph

Do you

$$y = \tan x$$
 and $y = -\cot\left(x + \frac{\pi}{2}\right)$
think that $\tan x = -\cot\left(x + \frac{\pi}{2}\right)$?

'Are You Prepared?' Answers


1.
$$x = 4$$
 2. True

5.6 Phase Shift; Sinusoidal Curve Fitting

OBJECTIVES 1 Graph Sinusoidal Functions of the Form $y = A \sin(\omega x - \phi)$, Using the Amplitude, Period, and Phase Shift

2 Find a Sinusoidal Function from Data

Figure 81 One cycle $y = A \sin(\omega x), A > 0, \omega > 0$



Graph Sinusoidal Functions of the Form $y = A \sin(\omega x - \phi)$, Using the Amplitude, Period, and Phase Shift

We have seen that the graph of $y = A \sin(\omega x)$, $\omega > 0$, has amplitude |A| and period $T = \frac{2\pi}{\omega}$. One cycle can be drawn as x varies from 0 to $\frac{2\pi}{\omega}$ or, equivalently, as ωx varies from 0 to 2π . See Figure 81.

We now want to discuss the graph of

$$y = A\sin(\omega x - \phi) = A\sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$$

where $\omega > 0$ and ϕ (the Greek letter phi) are real numbers. The graph will be a sine curve with amplitude |A|. As $\omega x - \phi$ varies from 0 to 2π , one period will be traced out. This period will begin when

$$\omega x - \phi = 0$$
 or $x = \frac{\phi}{\omega}$

and will end when

$$\omega x - \phi = 2\pi$$
 or $x = \frac{2\pi}{\omega} + \frac{\phi}{\omega}$

See Figure 82.

We see that the graph of $y = A \sin(\omega x - \phi) = A \sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$ is the same

as the graph of $y = A \sin(\omega x)$, except that it has been shifted $\frac{\phi}{\omega}$ units (to the right if $\phi > 0$ and to the left if $\phi < 0$). This number $\frac{\phi}{\omega}$ is called the **phase shift** of the graph of $y = A \sin(\omega x - \phi)$.

For the graphs of
$$y = A \sin(\omega x - \phi)$$
 or $y = A \cos(\omega x - \phi), \omega > 0$

Amplitude =
$$|A|$$
 Period = $T = \frac{2\pi}{\omega}$ Phase shift = $\frac{\phi}{\omega}$

The phase shift is to the left if $\phi < 0$ and to the right if $\phi > 0$.

EXAMPLE 1 Finding the Amplitude, Period, and Phase Shift of a Sinusoidal Function and Graphing It

Find the amplitude, period, and phase shift of $y = 3\sin(2x - \pi)$, and graph the function.


Solution Comparing

$$y = 3\sin(2x - \pi) = 3\sin\left[2\left(x - \frac{\pi}{2}\right)\right]$$

to

$$y = A\sin(\omega x - \phi) = A\sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$$

we find that A = 3, $\omega = 2$, and $\phi = \pi$. The graph is a sine curve with amplitude |A| = 3, period $T = \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi$, and phase shift $= \frac{\phi}{\omega} = \frac{\pi}{2}$.



One cycle $y = A \sin(\omega x - \phi), A > 0$,

Figure 82


 $\omega > 0, \phi > 0$

The graph of $y = 3\sin(2x - \pi)$ will lie between -3 and 3 on the y-axis. One cycle will begin at $x = \frac{\phi}{\omega} = \frac{\pi}{2}$ and end at $x = \frac{2\pi}{\omega} + \frac{\phi}{\omega} = \pi + \frac{\pi}{2} = \frac{3\pi}{2}$. We divide the interval $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ into four subintervals, each of length $\pi \div 4 = \frac{\pi}{4}$: $\left[\frac{\pi}{2}, \frac{3\pi}{4}\right], \left[\frac{3\pi}{4}, \pi\right], \left[\pi, \frac{5\pi}{4}\right], \left[\frac{5\pi}{4}, \frac{3\pi}{2}\right]$

The end points of these subintervals give rise to the following five key points on the graph:


$$\left(\frac{\pi}{2},0\right), \quad \left(\frac{3\pi}{4},3\right), \quad (\pi,0), \quad \left(\frac{5\pi}{4},-3\right), \quad \left(\frac{3\pi}{2},0\right)$$

We plot these five points and fill in the graph of the sine function as shown in Figure 83(a). Extending the graph in either direction, we obtain Figure 83(b).



✓ CHECK: Figure 85 shows the graph of $Y_1 = 3 \sin(2x - \pi)$ using a graphing utility.

Figure 85



EXAMPLE 2 Finding the Amplitude, Period, and Phase Shift of a Sinusoidal Function and Graphing It

Find the amplitude, period, and phase shift of $y = 2\cos(4x + 3\pi)$, and graph the function.

Solution Comparing

$$y = 2\cos(4x + 3\pi) = 2\cos\left[4\left(x + \frac{3\pi}{4}\right)\right]$$

to

$$y = A\cos(\omega x - \phi) = A\cos\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$$

we see that A = 2, $\omega = 4$, and $\phi = -3\pi$. The graph is a cosine curve with amplitude |A| = 2, period $T = \frac{2\pi}{\omega} = \frac{2\pi}{4} = \frac{\pi}{2}$, and phase shift $= \frac{\phi}{\omega} = -\frac{3\pi}{4}$.


The graph of $y = 2\cos(4x + 3\pi)$ will lie between -2 and 2 on the y-axis. One cycle will begin at $x = \frac{\phi}{\omega} = -\frac{3\pi}{4}$ and end at $x = \frac{2\pi}{\omega} + \frac{\phi}{\omega} = \frac{\pi}{2} + \left(-\frac{3\pi}{4}\right) = -\frac{\pi}{4}$. We divide the interval $\left[-\frac{3\pi}{4}, -\frac{\pi}{4}\right]$ into four subintervals, each of the length $\frac{\pi}{2} \div 4 = \frac{\pi}{8}$:

$$\left[-\frac{3\pi}{4},-\frac{5\pi}{8}\right], \quad \left[-\frac{5\pi}{8},-\frac{\pi}{2}\right], \quad \left[-\frac{\pi}{2},-\frac{3\pi}{8}\right], \quad \left[-\frac{3\pi}{8},-\frac{\pi}{4}\right]$$


The five key points on the graph are

$$\left(-\frac{3\pi}{4},2\right), \left(-\frac{5\pi}{8},0\right), \left(-\frac{\pi}{2},-2\right), \left(-\frac{3\pi}{8},0\right), \left(-\frac{\pi}{4},2\right)$$

We plot these five points and fill in the graph of the cosine function as shown in Figure 86(a). Extending the graph in either direction, we obtain Figure 86(b).



The graph of $y = 2\cos(4x + 3\pi) = 2\cos\left[4\left(x + \frac{3\pi}{4}\right)\right]$ may also be obtained using transformations. See Figure 87.



NOW WORK PROBLEM 3.

Summary

Steps for Graphing Sinusoidal Functions $y = A \sin(\omega x - \phi)$ or $y = A \cos(\omega x - \phi)$

STEP 1: Determine the amplitude |A| and period $T = \frac{2\pi}{\omega}$.

STEP 2: Determine the starting point of one cycle of the graph, $\frac{\phi}{\omega}$.

STEP 3: Determine the ending point of one cycle of the graph, $\frac{2\pi}{\omega} + \frac{\phi}{\omega}$.

STEP 4: Divide the interval $\left[\frac{\phi}{\omega}, \frac{2\pi}{\omega} + \frac{\phi}{\omega}\right]$ into four subintervals, each of length $\frac{2\pi}{\omega} \div 4$.

STEP 5: Use the endpoints of the subintervals to find the five key points on the graph.


STEP 6: Fill in one cycle of the graph.


STEP 7: Extend the graph in each direction to make it complete.

2 Find a Sinusoidal Function from Data

Scatter diagrams of data sometimes take the form of a sinusoidal function. Let's look at an example.

The data given in Table 12 represent the average monthly temperatures in Denver, Colorado. Since the data represent *average* monthly temperatures collected over many years, the data will not vary much from year to year and so will essentially repeat each year. In other words, the data are periodic. Figure 88 shows the scatter diagram of these data repeated over 2 years, where x = 1 represents January, x = 2 represents February, and so on.





•


25

$$y = A \sin(\omega x - \phi) + B$$

where A, B, ω , and ϕ are constants.

EXAMPLE 3 Finding a Sinusoidal Function from Temperature Data

Fit a sine function to the data in Table 12.



We begin with a scatter diagram of the data for 1 year. See Figure 89. The data will be fitted to a sine function of the form

$$y = A\sin(\omega x - \phi) + B$$

STEP 1: To find the amplitude *A*, we compute

Amplitude =
$$\frac{\text{largest data value} - \text{smallest data value}}{2}$$

= $\frac{73.5 - 29.7}{2} = 21.9$


To see the remaining steps in this process, we superimpose the graph of the function $y = 21.9 \sin x$, where x represents months, on the scatter diagram. Figure 90 shows the two graphs.

To fit the data, the graph needs to be shifted vertically, shifted horizontally, and stretched horizontally.

STEP 2: We determine the vertical shift by finding the average of the highest and lowest data value.

Vertical shift
$$=\frac{73.5 + 29.7}{2} = 51.6$$





Now we superimpose the graph of $y = 21.9 \sin x + 51.6$ on the scatter diagram. See Figure 91.

We see that the graph needs to be shifted horizontally and stretched horizontally.

STEP 3: It is easier to find the horizontal stretch factor first. Since the temperatures repeat every 12 months, the period of the function is T = 12. Since

$$T = \frac{2\pi}{\omega} = 12$$
, we have

$$\omega = \frac{2\pi}{12} = \frac{\pi}{6}$$

Now we superimpose the graph of $y = 21.9 \sin\left(\frac{\pi}{6}x\right) + 51.6$ on the scatter diagram. See Figure 92.

We see that the graph still needs to be shifted horizontally.

STEP 4: To determine the horizontal shift, we use the period T = 12 and divide the interval [0, 12] into four subintervals of length $12 \div 4 = 3$:

The sine curve is increasing on the interval (0, 3), and is decreasing on the interval (3, 9), so a local maximum occurs at x = 3. The data indicate that a maximum occurs at x = 7 (corresponding to July's temperature), so we must shift the graph of the function 4 units to the right by replacing x by x - 4. Doing this, we obtain

$$y = 21.9\sin\!\left(\frac{\pi}{6}(x-4)\right) + 51.6$$

Multiplying out, we find that a sine function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data is

$$y = 21.9 \sin\left(\frac{\pi}{6}x - \frac{2\pi}{3}\right) + 51.6$$

The graph of $y = 21.9 \sin\left(\frac{\pi}{6}x - \frac{2\pi}{3}\right) + 51.6$ and the scatter diagram of the data are shown in Figure 93.




Figure 92





Figure 93



The steps to fit a sine function

$$y = A\sin(\omega x - \phi) + B$$

to sinusoidal data follow:

Figure 91

Steps for Fitting Data to a Sine Function $y = A$ si	$m(\omega x - \phi) + B$
STEP 1: Determine <i>A</i> , the amplitude of the function.	
$Amplitude = \frac{largest data value - smal}{2}$	lest data value
STEP 2: Determine <i>B</i> , the vertical shift of the function.	
Vertical shift = $\frac{\text{largest data value + sma}}{2}$	llest data value
STEP 3: Determine ω . Since the period <i>T</i> , the time it takes peat, is $T = \frac{2\pi}{\omega}$, we have $\omega = \frac{2\pi}{T}$	kes for the data to re-
STEP 4: Determine the horizontal shift of the function the data. Divide the period into four subinter Determine the <i>x</i> -coordinate for the maximum and the <i>x</i> -coordinate for the maximum value information to determine the value of the phase	vals of equal length. of the sine function of the data. Use this
NOW WORK PROBLEMS 21(a)-(c).	

Let's look at another example. Since the number of hours of sunlight in a day cycles annually, the number of hours of sunlight in a day for a given location can be modeled by a sinusoidal function.

The longest day of the year (in terms of hours of sunlight) occurs on the day of the summer solstice. The summer solstice is the time when the sun is farthest north. In 2005, the summer solstice occurred on June 21 (the 172nd day of the year) at 2:46 AM EDT. The shortest day of the year occurs on the day of the winter solstice. The winter solstice is the time when the Sun is farthest south (again, for locations in the northern hemisphere). In 2005, the winter solstice occurred on December 21 (the 355th day of the year) at 1:35 PM (EST).

EXAMPLE 4 Finding a Sinusoidal Function for Hours of Daylight

According to the Old Farmer's Almanac, the number of hours of sunlight in Boston on the summer solstice is 15.283 and the number of hours of sunlight on the winter solstice is 9.067.

- (a) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the Old Farmer's Almanac and compare the actual hours of daylight to the results found in part (b).

Solution (a) STEP 1: Amplitude =
$$\frac{\text{largest data value} - \text{smallest data value}}{2}$$

= $\frac{15.283 - 9.067}{2}$ = 3.108
STEP 2: Vertical shift = $\frac{\text{largest data value} + \text{smallest data value}}{2}$
= $\frac{15.283 + 9.067}{2}$ = 12.175
STEP 3: The data repeat every 365 days. Since $T = \frac{2\pi}{\omega}$ = 365, we find
 $\omega = \frac{2\pi}{365}$
So far, we have $y = 3.108 \sin\left(\frac{2\pi}{365}x - \phi\right) + 12.175$.

STEP 4: To determine the horizontal shift, we use the period T = 365 and divide the interval [0, 365] into four subintervals of length $365 \div 4 = 91.25$:

$$[0, 91.25], [91.25, 182.5], [182.5, 273.75], [273.75, 365]$$

The sine curve is increasing on the interval (0, 91.25) and is decreasing on the interval (91.25, 273.75), so a local maximum occurs at x = 91.25. Since the maximum occurs on the summer solstice at x = 172, we must shift the graph of the function 172 - 91.25 = 80.75 units to the right by replacing x by x - 80.75. Doing this, we obtain

$$y = 3.108 \sin\left(\frac{2\pi}{365}(x - 80.75)\right) + 12.175$$

Multiplying out, we find that a sine function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data is

$$y = 3.108 \sin\left(\frac{2\pi}{365}x - \frac{323\pi}{730}\right) + 12.175$$

(b) To predict the number of hours of daylight on April 1, we let x = 91 in the function found in part (a) and obtain

$$y = 3.108 \sin\left(\frac{2\pi}{365} \cdot 91 - \frac{323}{730}\pi\right) + 12.175$$


\$\approx 12.72\$

So we predict that there will be about 12.72 hours = 12 hours, 43 minutes of sunlight on April 1 in Boston.

- (c) The graph of the function found in part (a) is given in Figure 94.
- (d) According to the *Old Farmer's Almanac*, there will be 12 hours 43 minutes of sunlight on April 1 in Boston. Our results agree with the Old Farmer's Almanac!


NOW WORK PROBLEM 27.

Figure 94



Certain graphing utilities (such as a TI-83, TI-84 Plus, and TI-86) have the capability of finding the sine function of best fit for sinusoidal data. At least four data points are required for this process.

EXAMPLE 5 Finding the Sine Function of Best Fit Use a graphing utility to find the sine function of best fit for the data in Table 12. Graph this function with the scatter diagram of the data. **Solution** Enter the data from Table 12 and execute the SINe REGression program. The result is shown in Figure 95. The output that the utility provides shows the equation $y = a\sin(bx + c) + d$ The sinusoidal function of best fit is $y = 21.15\sin(0.55x - 2.35) + 51.19$ where *x* represents the month and *y* represents the average temperature. Figure 96 shows the graph of the sinusoidal function of best fit on the scatter diagram. Figure 95 Figure 96 75



5.6 Assess Your Understanding

Concepts and Vocabulary

- **1.** For the graph of $y = A \sin(\omega x \phi)$, the number $\frac{\phi}{\omega}$ is called the _____.
- **2.** *True or False*: Only two data points are required by a graphing utility to find the sine function of best fit.

Skill Building

In Problems 3–14, find the amplitude, period, and phase shift of each function. Graph each function. Show at least one period. Verify the result using a graphing utility.

3. $y = 4\sin(2x - \pi)$	4. $y = 3\sin(3x - \pi)$	$5. \ y = 2\cos\left(3x + \frac{\pi}{2}\right)$
6. $y = 3\cos(2x + \pi)$	7. $y = -3\sin\left(2x + \frac{\pi}{2}\right)$	$8. \ y = -2\cos\left(2x - \frac{\pi}{2}\right)$
9. $y = 4\sin(\pi x + 2)$	10. $y = 2\cos(2\pi x + 4)$	11. $y = 3\cos(\pi x - 2)$
12. $y = 2\cos(2\pi x - 4)$	13. $y = 3\sin\left(-2x + \frac{\pi}{2}\right)$	14. $y = 3\cos\left(-2x + \frac{\pi}{2}\right)$

In Problems 15–18, write the equation of a sine function that has the given characteristics.

15. Amplitude: 2**16.** Amplitude: 3**17.** Amplitude: 3**18.** Amplitude: 2Period:
$$\pi$$
Period: $\frac{\pi}{2}$ Period: 3π Period: π Phase shift: $\frac{1}{2}$ Phase shift: 2Phase shift: $-\frac{1}{3}$ Phase shift: -2

Applications and Extensions

19. Alternating Current (ac) Circuits The current *I*, in amperes, flowing through an ac (alternating current) circuit at time *t* is

$$I = 120\sin\left(30\pi t - \frac{\pi}{3}\right), \qquad t \ge 0$$

What is the period? What is the amplitude? What is the phase shift? Graph this function over two periods.

20. Alternating Current (ac) Circuits The current *I*, in amperes, flowing through an ac (alternating current) circuit at time *t* is

$$I = 220\sin\left(60\pi t - \frac{\pi}{6}\right), \qquad t \ge 0$$

What is the period? What is the amplitude? What is the phase shift? Graph this function over two periods.

21. Monthly Temperature The following data represent the average monthly temperatures for Juneau, Alaska.

A CONTRACTOR	Month, x	Average Monthly Temperature, °F
7.0	January, 1	24.2
· · · ·	February, 2	28.4
	March, 3	32.7
	April, 4	39.7
	May, 5	47.0
	June, 6	53.0
	July, 7	56.0
	August, 8	55.0
	September, 9	49.4
	October, 10	42.2
	November, 11	32.0
	December, 12	27.1

SOURCE: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Draw the sinusoidal function of best fit on the scatter diagram.

22. Monthly Temperature The following data represent the average monthly temperatures for Washington, D.C.

Month, <i>x</i>	Average Monthly Temperature, °F
January, 1	34.6
February, 2	37.5
March, 3	47.2
April, 4	56.5
May, 5	66.4
June, 6	75.6
July, 7	80.0
August, 8	78.5
September, 9	71.3
October, 10	59.7
November, 11	49.8
December, 12	39.4

Source: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on the scatter diagram.
- **23. Monthly Temperature** The following data represent the average monthly temperatures for Indianapolis, Indiana.

	Month, <i>x</i>	Average Monthly Temperature, °F
1~	January, 1	25.5
	February, 2	29.6
	March, 3	41.4
	April, 4	52.4
	May, 5	62.8
	June, 6	71.9
	July, 7	75.4
	August, 8	73.2
	September, 9	66.6
	October, 10	54.7
	November, 11	43.0
	December, 12	30.9

Source: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on the scatter diagram.
- **24. Monthly Temperature** The following data represent the average monthly temperatures for Baltimore, Maryland.

The second second	Month, <i>x</i>	Average Monthly Temperature, °F
	January, 1	31.8
	February, 2	34.8
	March, 3	44.1
	April, 4	53.4
	May, 5	63.4
	June, 6	72.5
	July, 7	77.0
	August, 8	75.6
	September, 9	68.5
	October, 10	56.6
	November, 11	46.8
	December, 12	36.7

Source: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on the scatter diagram.
- **25. Tides** Suppose that the length of time between consecutive high tides is approximately 12.5 hours. According to the National Oceanic and Atmospheric Administration, on Saturday, August 7, 2004, in Savannah, Georgia, high tide occurred at 3:38 AM (3.6333 hours) and low tide occurred at 10:08 AM (10.1333 hours). Water heights are measured as the amounts above or below the mean lower low water. The height of the water at high tide was 8.2 feet, and the height of the water at low tide was -0.6 foot.
 - (a) Approximately when will the next high tide occur?
 - (b) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
 - (c) Draw a graph of the function found in part (b).
 - (d) Use the function found in part (b) to predict the height of the water at the next high tide.

- **26. Tides** Suppose that the length of time between consecutive high tides is approximately 12.5 hours. According to the National Oceanic and Atmospheric Administration, on Saturday, August 7, 2004, in Juneau, Alaska, high tide occurred at 8:11 AM (8.1833 hours) and low tide occurred at 2:14 PM (14.2333 hours). Water heights are measured as the amounts above or below the mean lower low water. The height of the water at high tide was 13.2 feet, and the height of the water at low tide was 2.2 feet.
 - (a) Approximately when will the next high tide occur?
 - (b) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
 - (c) Draw a graph of the function found in part (b).
 - (d) Use the function found in part (b) to predict the height of the water at the next high tide.
- 27. Hours of Daylight According to the *Old Farmer's Almanac*, in Miami, Florida, the number of hours of sunlight on the summer solstice is 12.75 and the number of hours of sunlight on the winter solstice is 10.583.
 - (a) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
 - (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
 - (c) Draw a graph of the function found in part (a).
 - (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
 - **28. Hours of Daylight** According to the *Old Farmer's Almanac*, in Detroit, Michigan, the number of hours of sunlight on the summer solstice is 13.65 and the number of hours of sunlight on the winter solstice is 9.067.
 - (a) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
 - (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
 - (c) Draw a graph of the function found in part (a).
 - (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
 - **29. Hours of Daylight** According to the *Old Farmer's Almanac*, in Anchorage, Alaska, the number of hours of sunlight on the summer solstice is 16.233 and the number of hours of sunlight on the winter solstice is 5.45.
 - (a) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
 - (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
 - (c) Draw a graph of the function found in part (a).
 - (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
 - **30. Hours of Daylight** According to the *Old Farmer's Almanac*, in Honolulu, Hawaii, the number of hours of sunlight on the summer solstice is 12.767 and the number of hours of sunlight on the winter solstice is 10.783.

- (a) Find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.

Discussion and Writing

31. Explain how the amplitude and period of a sinusoidal graph are used to establish the scale on each coordinate axis.

- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
- **32.** Find an application in your major field that leads to a sinusoidal graph. Write a paper about your findings.

Chapter Review

Things to Know

Definitions

Angle in standard position (p. 356)	Vertex is at the origin; initial side is along the positive <i>x</i> -axis		
1 Degree (1°) (p. 357)	$1^\circ = \frac{1}{360}$ revolution		
1 Radian (p. 360)	The measure of a central angle of a circle whose rays subtend an arc whose length is the radius of the circle		
Trigonometric functions (pp. 371–372)	$P = (x, y)$ is the point on the unit circle corresponding to $\theta = t$ radians.		
	$\sin t = \sin \theta = y$ $\cos t = \cos \theta = x$ $\tan t = \tan \theta = \frac{y}{x}, x \neq 0$		
	$\csc t = \csc \theta = \frac{1}{y}, y \neq 0$ $\sec t = \sec \theta = \frac{1}{x}, x \neq 0$ $\cot t = \cot \theta = \frac{x}{y}, y \neq 0$		
Trigonometric functions using a circle of radius r (pp. 382–383)	For an angle θ in standard position $P = (x, y)$ is the point on the terminal side of θ that is also on the circle $x^2 + y^2 = r^2$.		
	$\sin\theta = \frac{y}{r}$ $\cos\theta = \frac{x}{r}$ $\tan\theta = \frac{y}{x}, x \neq 0$		
	$\csc \theta = \frac{r}{y}, y \neq 0$ $\sec \theta = \frac{r}{x}, x \neq 0$ $\cot \theta = \frac{x}{y}, y \neq 0$		
Periodic function (p. 391)	$f(\theta + p) = f(\theta)$, for all θ , $p > 0$, where the smallest such p is the fundamental period		
Formulas			
1 revolution = 360° (p. 358) = 2π radians (p. 361)			
$s = r \theta \text{ (p. 360)}$	θ is measured in radians; s is the length of arc subtended by the central angle θ of the circle of radius r; A is the area of the sector.		
$A = \frac{1}{2}r^2\theta \text{ (p. 364)}$			

 $v = r\omega$ (p. 365)

v is the linear speed along the circle of radius r; ω is the angular speed (measured in radians per unit time).

TABLE OF VALUES							
θ (Radians)	θ (Degrees)	$\sin heta$	$\cos heta$	tan θ	$\csc \theta$	$\sec \theta$	$\cot heta$
0	0°	0	1	0	Not defined	1	Not defined
$\frac{\pi}{6}$	30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
$\frac{\pi}{4}$	45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
$\frac{\pi}{3}$	60°	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$
$\frac{\pi}{2}$	90°	1	0	Not defined	1	Not defined	0
π	180°	0	-1	0	Not defined	-1	Not defined
$\frac{3\pi}{2}$	270°	-1	0	Not defined	-1	Not defined	0

 θ


Fundamental Identities (p. 394)


$$\tan \theta = \frac{\sin \theta}{\cos \theta}, \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$
$$\csc \theta = \frac{1}{\sin \theta}, \quad \sec \theta = \frac{1}{\cos \theta}, \quad \cot \theta = \frac{1}{\tan \theta}$$
$$\sin^2 \theta + \cos^2 \theta = 1, \quad \tan^2 \theta + 1 = \sec^2 \theta, \quad 1 + \cot^2 \theta = \csc^2 \theta$$


Properties of the Trigonometric Functions


- $y = \sin x \qquad \text{Domain:} -\infty < x < \infty$ (p. 404) $\begin{array}{l} \text{Range:} -1 \le y \le 1 \\ \text{Periodic: period} = 2\pi (360^\circ) \\ \text{Odd function} \end{array}$
- y = cos x Domain: $-\infty < x < \infty$ (p. 406) Range: $-1 \le y \le 1$ Periodic: period = 2π (360°) Even function
- $y = \tan x \qquad \text{Domain:} -\infty < x < \infty, \text{ except odd multiples of } \frac{\pi}{2}(90^\circ)$ (p. 420) $\begin{array}{l} \text{Range:} -\infty < y < \infty \\ \text{Periodic: period} = \pi (180^\circ) \\ \text{Odd function} \end{array}$

y = cot x Domain: $-\infty < x < \infty$, except integer multiples of π (180°) (p. 422) Range: $-\infty < y < \infty$ Periodic: period = π (180°) Odd function










y = csc x Domain: $-\infty < x < \infty$, except integer multiples of π (180°) (p. 422) Range: $|y| \ge 1$ Periodic: period = 2π (360°) Odd function



y = sec x Domain: $-\infty < x < \infty$, except odd multiples of $\frac{\pi}{2}(90^{\circ})$ (p. 423) Range: $|y| \ge 1$ Periodic: period = 2π (360°) Even function

Sinusoidal graphs (pp. 409 and 426)

$y = A\sin(\omega x), \omega > 0$	Period = $\frac{2\pi}{\omega}$
$y = A\cos(\omega x), \omega > 0$	Amplitude = $ A $
$y = A\sin(\omega x - \phi) = A\sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$	Phase shift $= \frac{\phi}{\omega}$
$y = A\cos(\omega x - \phi) = A\cos\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$	

Objectives

Secti	on	You should be able to	Review Exercises
5.1	1	Convert between degrees, minutes, seconds, and decimal forms for angles (p. 358)	82
	2	Find the arc length of a circle (p. 360)	83,84
	3	Convert from degrees to radians and from radians to degrees (p. 361)	1-8
	4	Find the area of a sector of a circle (p. 364)	83
	5	Find the linear speed of an object traveling in circular motion (p. 365)	85–88
5.2	J	Find the exact values of the trigonometric functions using a point on the unit circle (p. 372)	79
	2	Find the exact values of the trigonometric functions of quadrantal angles (p. 374)	17, 18, 20
	3	Find the exact values of the trigonometric functions of $\frac{\pi}{4} = 45^{\circ}$ (p. 376)	9, 11, 13, 15, 16, 19
	4	Find the exact values of the trigonometric functions of $\frac{\pi}{6} = 30^{\circ}$ and $\frac{\pi}{3} = 60^{\circ}$ (p. 377)	9–15
	5	Find the exact values of the trigonometric functions for integer multiples of $\frac{\pi}{6} = 30^{\circ}, \frac{\pi}{4} = 45^{\circ},$	
		and $\frac{\pi}{3} = 60^{\circ}$ (p. 380)	13–16,94
	6	Use a calculator to approximate the value of a trigonometric function (p. 381)	75,76
	J	Use circle of radius r to evaluate the trigonometric functions (p. 388)	80
5.3	J	Determine the domain and the range of the trigonometric functions (p. 388)	81
	2	Determine the period of the trigonometric functions (p. 390)	81
	3	Determine the signs of the trigonometric functions in a given quadrant (p. 392)	77–78
	4	Find the values of the trigonometric functions using fundamental identities (p. 393)	21–30
	5	Find the exact values of the trigonometric functions of an angle given one of the functions and the quadrant of the angle (p. 395)	31–46

440 CHAPTER 5 Trigonometric Functions

	6	Use even-odd properties to find the exact values of the trigonometric functions (p. 398)	27–30
5.4	1	Graph transformations of the sine function (p. 403)	47, 50
	2	Graph transformations of the cosine function (p. 405)	48, 49
	3	Determine the amplitude and period of sinusoidal functions (p. 408)	59-64, 89
	4	Graph sinusoidal functions using key points: (p. 410)	47, 48, 63-64, 89
	5	Find an equation for a sinusoidal graph (p. 413)	71–74
5.5	1	Graph transformations of the tangent function and cotangent function (p. 419)	51–56
	2	Graph transformations of the cosecant function and secant function (p. 422)	57, 58
5.6	3	Graph sinusoidal functions of the form $y = A \sin(\omega x - \phi)$ using the amplitude, period and phase shift (p. 425)	65–70, 90
	2	Find a sinusoidal function from data (p. 429)	91–93

Review Exercises

In Problems 1–4, convert each angle in degrees to radians. Express your answer as a multiple of π .

1. 135° 2. 210° 3. 18°	4. 15°
---	---------------

In Problems 5-8, convert each angle in radians to degrees.

5.
$$\frac{3\pi}{4}$$
 6. $\frac{2\pi}{3}$ **7.** $-\frac{5\pi}{2}$ **8.** $-\frac{3\pi}{2}$

In Problems 9–30, find the exact value of each expression. Do not use a calculator.

10. $\cos \frac{\pi}{3} + \sin \frac{\pi}{2}$ 9. $tan \frac{\pi}{4} - sin \frac{\pi}{6}$ **11.** $3\sin 45^\circ - 4\tan \frac{\pi}{6}$ **13.** $6\cos\frac{3\pi}{4} + 2\tan\left(-\frac{\pi}{3}\right)$ **14.** $3\sin\frac{2\pi}{3} - 4\cos\frac{5\pi}{2}$ **12.** $4\cos 60^\circ + 3\tan \frac{\pi}{3}$ 15. $\operatorname{sec}\left(-\frac{\pi}{3}\right) - \operatorname{cot}\left(-\frac{5\pi}{4}\right)$ 16. $4 \csc \frac{3\pi}{4} - \cot \left(-\frac{\pi}{4}\right)$ 17. $\tan \pi + \sin \pi$ 18. $\cos\frac{\pi}{2} - \csc\left(-\frac{\pi}{2}\right)$ **19.** $\cos 540^\circ - \tan(-45^\circ)$ **20.** $\sin 630^\circ + \cos(-180^\circ)$ **21.** $\sin^2 20^\circ + \frac{1}{\sec^2 20^\circ}$ **22.** $\frac{1}{\cos^2 40^\circ} - \frac{1}{\cot^2 40^\circ}$ **23.** sec 50° cos 50° **26.** $\frac{1}{\sec^2 40^\circ} + \frac{1}{\csc^2 40^\circ}$ **25.** $\sec^2 20^\circ - \tan^2 20^\circ$ **24.** tan 10° cot 10° **27.** $\sin(-40^\circ) \csc 40^\circ$ **28.** $tan(-20^\circ) \cot 20^\circ$ **29.** $\cos 410^{\circ} \sec(-50^{\circ})$ **30.** $\cot 200^{\circ} \tan(-20^{\circ})$

In Problems 31–46, find the exact value of each of the remaining trigonometric functions.

31.
$$\sin \theta = \frac{4}{5}$$
, θ is acute
32. $\cos \theta = \frac{3}{5}$, θ is acute
33. $\tan \theta = \frac{12}{5}$, $\sin \theta < 0$
34. $\cot \theta = \frac{12}{5}$, $\cos \theta < 0$
35. $\sec \theta = -\frac{5}{4}$, $\tan \theta < 0$
36. $\csc \theta = -\frac{5}{3}$, $\cot \theta < 0$

Chapter Review 441

37.
$$\sin \theta = \frac{12}{13}$$
, θ in quadrant II
38. $\cos \theta = -\frac{3}{5}$, θ in quadrant III
39. $\sin \theta = -\frac{5}{13}$, $\frac{3\pi}{2} < \theta < 2\pi$
40. $\cos \theta = \frac{12}{13}$, $\frac{3\pi}{2} < \theta < 2\pi$
41. $\tan \theta = \frac{1}{3}$, $180^{\circ} < \theta < 270^{\circ}$
42. $\tan \theta = -\frac{2}{3}$, $90^{\circ} < \theta < 180^{\circ}$
43. $\sec \theta = 3$, $\frac{3\pi}{2} < \theta < 2\pi$
44. $\csc \theta = -4$, $\pi < \theta < \frac{3\pi}{2}$
45. $\cot \theta = -2$, $\frac{\pi}{2} < \theta < \pi$
46. $\tan \theta = -2$, $\frac{3\pi}{2} < \theta < 2\pi$

In Problems 47–58, graph each function. Each graph should contain at least one period.

47.
$$y = 2\sin(4x)$$
48. $y = -3\cos(2x)$ **49.** $y = -2\cos\left(x + \frac{\pi}{2}\right)$ **50.** $y = 3\sin(x - \pi)$ **51.** $y = \tan(x + \pi)$ **52.** $y = -\tan\left(x - \frac{\pi}{2}\right)$ **53.** $y = -2\tan(3x)$ **54.** $y = 4\tan(2x)$ **55.** $y = \cot\left(x + \frac{\pi}{4}\right)$ **56.** $y = -4\cot(2x)$ **57.** $y = \sec\left(x - \frac{\pi}{4}\right)$ **58.** $y = \csc\left(x + \frac{\pi}{4}\right)$

In Problems 59–62, determine the amplitude and period of each function without graphing.

59.
$$y = 4 \cos x$$
 60. $y = \sin(2x)$ **61.** $y = -8 \sin\left(\frac{\pi}{2}x\right)$ **62.** $y = -2\cos(3\pi x)$

In Problems 63–70, find the amplitude, period, and phase shift of each function. Graph each function. Show at least one period.


63.
$$y = 4\sin(3x)$$

64. $y = 2\cos\left(\frac{1}{3}x\right)$
65. $y = 2\sin(2x - \pi)$
66. $y = -\cos\left(\frac{1}{2}x + \frac{\pi}{2}\right)$
67. $y = \frac{1}{2}\sin\left(\frac{3}{2}x - \pi\right)$
68. $y = \frac{3}{2}\cos(6x + 3\pi)$
69. $y = -\frac{2}{3}\cos(\pi x - 6)$
70. $y = -7\sin\left(\frac{\pi}{3}x + \frac{4}{3}\right)$

73.

In Problems 71–74, find a function whose graph is given.

71.



72.

- $\begin{array}{c} y_{1} \\ 6 \\ -4 \\ -4 \\ -6 \end{array}$
- **75.** Use a calculator to approximate $\sin \frac{\pi}{8}$. Round the answer to two decimal places.
- **76.** Use a calculator to approximate sec 10°. Round the answer to two decimal places.
- **77.** Determine the signs of the six trigonometric functions of an angle θ whose terminal side is in quadrant III.
- **78.** Name the quadrant θ lies in if $\cos \theta > 0$ and $\tan \theta < 0$.
- $P = \left(-\frac{1}{3}, \frac{2\sqrt{2}}{3}\right)$ is the point on the unit circle that corresponds to *t*.

79. Find the exact values of the six trigonometric functions if

74.

80. Find the exact value of sin *t*, cos *t*, and tan *t* if P = (-2, 5) is the point on the circle that corresponds to *t*.

442 CHAPTER 5 Trigonometric Functions

- **81.** What is the domain and the range of the secant function? What is the period?
- **82.** (a) Convert the angle 32°20′35″ to a decimal in degrees. Round the answer to two decimal places.
 - (b) Convert the angle 63.18° to D°M'S" form. Express the answer to the nearest second.
- **83.** Find the length of the arc subtended by a central angle of 30° on a circle of radius 2 feet. What is the area of the sector?
- **84.** The minute hand of a clock is 8 inches long. How far does the tip of the minute hand move in 30 minutes? How far does it move in 20 minutes?
- **85. Angular Speed of a Race Car** A race car is driven around a circular track at a constant speed of 180 miles

per hour. If the diameter of the track is $\frac{1}{2}$ mile, what is

the angular speed of the car? Express your answer in revolutions per hour (which is equivalent to laps per hour).

- **86. Merry-Go-Rounds** A neighborhood carnival has a merrygo-round whose radius is 25 feet. If the time for one revolution is 30 seconds, how fast is the merry-go-round going?
- **87. Lighthouse Beacons** The Montauk Point Lighthouse on Long Island has dual beams (two light sources opposite each other). Ships at sea observe a blinking light every 5 seconds. What rotation speed is required to do this?
- **88. Spin Balancing Tires** The radius of each wheel of a car is 16 inches. At how many revolutions per minute should a spin balancer be set to balance the tires at a speed of 90 miles per hour? Is the setting different for a wheel of radius 14 inches? If so, what is this setting?
- **89.** Alternating Voltage The electromotive force *E*, in volts, in a certain ac (alternating circuit) circuit obeys the equation

$$E = 120\sin(120\pi t), \qquad t \ge 0$$

where *t* is measured in seconds.

- (a) What is the maximum value of *E*?
- (b) What is the period?
- (c) Graph this function over two periods.
- **90. Alternating Current** The current *I*, in amperes, flowing through an ac (alternating current) circuit at time *t* is

$$I = 220\sin\left(30\pi t + \frac{\pi}{6}\right), \qquad t \ge 0$$

- (a) What is the period?
- (b) What is the amplitude?
- (c) What is the phase shift?
- (d) Graph this function over two periods.
- **91. Monthly Temperature** The following data represent the average monthly temperatures for Phoenix, Arizona.

Month, <i>m</i>	Average Monthly Temperature, T
January, 1	51
February, 2	55
March, 3	63
April, 4	67
May, 5	77
June, 6	86
July, 7	90
August, 8	90
September, 9	84
October, 10	71
November, 11	59
December, 12	52

SOURCE: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on the scatter diagram.
- **92. Monthly Temperature** The following data represent the average monthly temperatures for Chicago, Illinois.

	Month, <i>m</i>	Average Monthly Temperature, T
\mathbf{r}	January, 1	25
J.	February, 2	28
	March, 3	36
	April, 4	48
	May, 5	61
	June, 6	72
	July, 7	74
	August, 8	75
	September, 9	66
	October, 10	55
	November, 11	39
	December, 12	28

SOURCE: U.S. National Oceanic and Atmospheric Administration

- (a) Use a graphing utility to draw a scatter diagram of the data for one period.
- (b) By hand, find a sinusoidal function of the form $y = A \sin(\omega x \phi) + B$ that fits the data.


- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
- (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on the scatter diagram.
- 93. Hours of Daylight According to the Old Farmer's Almanac, in Las Vegas, Nevada, the number of hours of sunlight on the summer solstice is 13.367 and the number of hours of sunlight on the winter solstice is 9.667. (a) Find a sinusoidal function of the form

$$y = A\sin(\omega x - \phi) + B$$

that fits the data.

- (b) Draw a graph of the function found in part (a).
- (c) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (d) Look up the number of hours of sunlight for April 1 in the Old Farmer's Almanac and compare the actual hours of daylight to the results found in part (c).

94. Unit Circle Fill in the angles (in degrees and radians) and terminal points P of each angle on the unit circle shown.



4

18. If $f(x) = \sin x$ and $f(a) = \frac{3}{5}$, find f(-a).

Chapter Test

In Problems 1–3, co	onvert each angle in degrees to radians. Express your answer as	s a multiple of π .
1. 260°	2. -400°	3. 13°

In Problems 4-6 convert each angle in radians to degrees.

4.
$$-\frac{\pi}{8}$$
 5. $\frac{9\pi}{2}$ **6.** $\frac{3\pi}{4}$

In Problems 7–12, find the exact value of each expression.

7.
$$\sin \frac{\pi}{6}$$
8. $\cos\left(-\frac{5\pi}{4}\right) - \cos \frac{3\pi}{4}$ 9. $\cos(-120^{\circ})$ 10. $\tan 330^{\circ}$ 11. $\sin \frac{\pi}{2} - \tan \frac{19\pi}{4}$ 12. $2\sin^2 60^{\circ} - 3\cos 45^{\circ}$

In Problems 13–16, use a calculator to evaluate each expression. Round your answers to three decimal places.

14. $\cos \frac{2\pi}{5}$ **16.** $\cot \frac{28\pi}{9}$ **15.** sec 229° **13.** sin 17°

17. Fill in each table entry with the sign of each function.

	$\sin \theta$	$\cos \theta$	tan θ	$\sec \theta$	$\csc \theta$	$\cot heta$
$\boldsymbol{\theta}$ in QI						
θ in QII						
$\boldsymbol{\theta}$ in QIII						
θ in QIV						

444 CHAPTER 5 Trigonometric Functions

In Problems 19–21 find the value of the remaining five trigonometric functions of θ .

19.
$$\sin \theta = \frac{5}{7}$$
, θ in quadrant II
20. $\cos \theta = \frac{2}{3}$, $\frac{3\pi}{2} < \theta < 2\pi$
21. $\tan \theta = -\frac{12}{5}$, $\frac{\pi}{2} < \theta < \pi$

In Problems 22–24, the point (x, y) is on the terminal side of angle θ in standard position. Find the exact value of the given trigonometric function.

22. (2, 7), sin θ

23. $(-5, 11), \cos \theta$

24. (6, -3), tan θ

In Problems 25 and 26, graph the function by hand.

 $25. \ y = 2\sin\left(x - \frac{\pi}{6}\right)$

26. $y = tan\left(-x + \frac{\pi}{4}\right) + 2$

27. Write an equation for a sinusoidal graph with the following properties:

$$A = -3$$
 period $= \frac{2\pi}{3}$ phase shift $= -\frac{\pi}{4}$


28. Logan has a garden in the shape of a sector of a circle; the outer rim of the garden is 25 ft long and the central angle of the sector is 50°. She wants to add a 3 ft wide walk to the outer rim; how many square feet of paving blocks will she need to build the walk?

29. Hungarian Adrian Annus won the gold medal for the hammer throw at the 2004 Olympics in Athens with a winning distance of 83.19 meters.^{*} The event consists of swinging a 16 lb weight—attached to a wire 190 cm long—in a circle, then releasing it. Assuming his release is at a 45° angle to the ground,

the hammer will travel a distance of $\frac{v_0^2}{g}$ meters, where $g = 9.8 \text{ m/sec}^2$ and v_0 is the linear speed of the hammer when

released. At what rate (rpm) was he swinging the hammer upon release?

- 30. The Freedom Tower is to be the centerpiece of the rebuilding of the World Trade Center in New York City. The tower will be 1776 feet tall (not including a broadcast antenna). The angle of elevation from the base of an office building to the top of the tower is 34°. The angle of elevation from the helipad on the roof of the office building to the top of the tower is 20°.
 - (a) How far away is the office building from the Freedom Tower (assume the side of the tower is vertical)? Round to the nearest foot.
 - (b) How tall is the office building? Round to the nearest foot.



*Annus was stripped of his medal after refusing post-medal drug testing.

Chapter Projects



- 1. **Tides** A partial tide table for September 2001 for Sabine Pass along the Texas Gulf Coast is given in the table.
 - (a) On September 15, when was the tide high? This is called *high tide*. On September 19, when was the tide low? This is called *low tide*. Most days will have two low tides and two high tides.
 - (b) Why do you think there is a negative height for the low tide on September 14? What is the tide height measured against?

- (c) On your graphing utility, draw a scatter diagram for the data in the table. Let T (time) be the independent variable, with T = 0 being 12:00 AM on September 1, T = 24 being 12:00 AM on September 2, and so on. Remember that there are 60 minutes in an hour. Let H be the height in feet. Also, make sure that your graphing utility is in radian mode.
- (d) What shape does the data take? What is the period of the data? What is the amplitude? Is the amplitude constant? Explain.
- (e) Using Steps 1–4 given on page 432, fit a sine curve to the data. Let the amplitude be the average of the amplitudes that you found in part (c), unless the amplitude was constant. Is there a vertical shift? Is there a phase shift?
- (f) Using your graphing utility, find the sinusoidal function of best fit. How does it compare to your equation?
- (g) Using the equation found in part (e) and the sinusoidal equation of best fit found in part (f), predict the high tides and the low tides on September 21.
- (h) Looking at the times of day that the low tides occur, what do you think causes the low tides to vary so much each day? Explain. Does this seem to have the same type of effect on the high tides? Explain.

Sept	High Time	Tide Ht (ft)	High Time	Tide Ht (ft)	Low Time	Tide Ht (ft)	Low Time	Tide Ht (ft)	Sun/Moon phase Rise/Set
14	03:08a	2.4	11:12a	2.2	08:14a	2.0	07:19p	-0.1	7:00a/7:23p
15	03:33a	2.4	12:56р	2.2	08:15a	1.9	08:13p	0.0	7:00a/7:22p
16	03:57a	2.3	02:17 _P	2.3	08:45a	1.6	09:05p	0.3	7:01a/7:20p
17	04:20a	2.2	03:33p	2.3	09:24 a	1.4	09:54p	0.5	7:01a/7:19p
18	04:41a	2.2	04:47 _P	2.3	10:08a	1.0	10:43p	1.0	7:02a/7:08p
19	05:01a	2.0	06:04p	2.3	10:54a	0.7	11:32p	1.4	7:02a/7:17p
20	05:20a	2.0	07:27p	2.3	11:44a	0.4			7:03a/7:15p


SOURCE: www.harbortides.com

The following projects are available at the Instructor's Resource Center (IRC):


- 2. Project at Motorola Digital Transmission Over the Air
- 3. Identifying Mountain Peaks in Hawaii
- 4. CBL Experiment

Cumulative Review

- **1.** Find the real solutions, if any, of the equation $2x^2 + x 1 = 0$.
- **2.** Find an equation for the line with slope -3 containing the point (-2, 5).
- **3.** Find an equation for a circle of radius 4 and center at the point (0, -2).
- 4. Discuss the equation 2x 3y = 12. Graph it.
- 5. Discuss the equation $x^2 + y^2 2x + 4y 4 = 0$. Graph it.
- 6. Use transformations to graph the function $y = (x - 3)^2 + 2.$
- **7.** Sketch a graph of each of the following functions. Label at least three points on each graph.
 - (a) $y = x^{2}$ (b) $y = x^{3}$ (c) $y = e^{x}$ (d) $y = \ln x$ (e) $y = \sin x$ (f) $y = \tan x$
- 8. Find the inverse function of f(x) = 3x 2.
- 9. Find the exact value of $(\sin 14^\circ)^2 + (\cos 14^\circ)^2 3$.
- **10.** Graph $y = 3 \sin(2x)$.
- **11.** Find the exact value of $\tan \frac{\pi}{4} 3\cos \frac{\pi}{6} + \csc \frac{\pi}{6}$.
- 12. Find an exponential function for the following graph. Express your answer in the form $y = Ab^x$.



13. Find a sinusoidal function for the following graph.



- 14. (a) Find a linear function that contains the points (-2, 3) and (1, -6). What is the slope? What are the intercepts of the function? Graph the function. Be sure to label the intercepts.
 - (b) Find a quadratic function that contains the point (−2, 3) with vertex (1, −6). What are the intercepts of the function? Graph the function.
 - (c) Show that there is no exponential function of the form $f(x) = ae^x$ that contains the points (-2, 3) and (1, -6).
- **15.** (a) Find a polynomial function of degree 3 whose *y*-intercept is 5 and whose *x*-intercepts are -2, 3, and 5. Graph the function. Label the local minima and local maxima.
 - (b) Find a rational function whose y-intercept is 5 and whose x-intercepts are -2, 3, and 5 that has the line x = 2 as a vertical asymptote. Graph the function. Answers may vary.